

Preplanned Studies

Weather Variability, Socioeconomic Factors, and Pneumonia in Children Under Five-Years Old — Bangladesh, 2012–2016

Mohammad Zahid Hossain^{1,2}; Shilu Tong^{1,3,4,5}; Hilary Bambrick¹; Md Alfazal Khan⁶; Wenbiao Hu^{1,*}

Summary

What is already known on this topic?

Different socioecological factors were associated with childhood pneumonia in Bangladesh. However, previous studies did not assess spatial patterns, and socioecological factors and spatial variation have the potential to improve the accuracy and predictive ability of existing models.

What is added by this report?

The spatial random effects were present at the district level and were heterogeneous. Average temperature, temperature variation, and population density may influence the spatial pattern of childhood pneumonia in Bangladesh.

What are the implications for public health practice?

The study results will help policymakers and health managers to identify the vulnerable districts, plan further investigations, help to improve proper resource allocation, and improve health interventions.

Pneumonia is one of the leading causes of mortality and morbidity in children aged under five years in Bangladesh. This study aimed to identify the association between weather, social factors and childhood pneumonia and identify the spatial variation of the disease. A Bayesian spatial Poisson regression model with a conditional autoregressive prior structure was developed to quantify the association between childhood pneumonia and socioecological factors and identify the spatial variation. The study results suggested that a 1 °C increase in monthly temperature and monthly temperature variation may increase the monthly associated log relative risk (RR) of childhood pneumonia by 1.161 [95% credible interval (CrI): 1.013–1.429] and 1.463 (95% CrI: 1.170–1.839), respectively. However, the population density was inversely related with pneumonia risk (RR: 0.996, CrI: 0.994–0.998). Socioecological factors may influence the spatial pattern of childhood pneumonia, and the spatial random effects were heterogeneous.

The study was conducted in Bangladesh, which is located in the northeastern part of South Asia. Bangladesh is divided into 8 administrative divisions and 64 districts. Monthly data on under-5-years pneumonia were extracted from the District Health Information System Version 2 of the Directorate General of Health Services (DGHS) under the Ministry of Health and Family Welfare of Bangladesh from January 2012 to December 2016 (1). The pneumonia cases were diagnosed according to the World Health Organization pneumonia guidelines (2). The under-five-years population data at the district level were collected from the latest national population and household census (3). The sociodemographic data (percentage of education and internet use) at the district level were collected from socioeconomic and demographic reports (national series, volume-4) from the same census. The poverty data for each district was obtained from the Household Income and Expenditure Survey 2016 (4).

Climate data (temperature and rainfall) were obtained from the National Environmental Satellite, Data and Information service (<https://www7.ncdc.noaa.gov/CDO/cdoselect.cmd?datasetabbv=GSOD>), which is publicly available and widely used in previous studies (5–6). Poisson regression models in a Bayesian framework were developed for pneumonia cases at the district level. These models assume that the observed counts of childhood pneumonia cases (O_k) for the k th district ($k=1\cdots 64$) follow a Poisson distribution with mean μ_k :

$$O_k \sim \text{Poisson}(\mu_k) \quad (1)$$

$$\log(\mu_k) = \log(E_k) + \theta_k \quad (2)$$

where E_k (the expected number of cases in District _{k}) is an offset to control population size and θ_k is the associated log RR.

Prior to this analysis, we examined multicollinearity among the different covariates but did not find sufficiently strong associations to warrant exclusion or other treatment of any variables (Supplementary Table S1, available in <http://weekly.chinacdc.cn>). As a consequence, a total of 6 models were developed

(Supplementary Material, available in <http://weekly.chinacdc.cn>). The model which incorporated all socioecological covariates with both structured and unstructured random effects were selected for the final analysis.

The expected log relative risk θ_k was represented as follows:

$$\begin{aligned}\theta_k = & \alpha + (\text{Temp}_k) \beta_1 + (\text{Tempva}_k) \beta_2 + (\text{Rain}_k) \beta_3 \\ & + (\text{Edu}_k) \beta_4 + (\text{Int}_k) \beta_5 + (\text{povi}_k) \beta_6 \\ & + (\text{pop}_k) \beta_7 + u_k + v_k\end{aligned}$$

where α is a constant; β_1 is the coefficient for temperature, β_2 is the coefficient for temperature variation, β_3 is the coefficient for rainfall, β_4 is the coefficient for percentage of education at the district level, β_5 is the coefficient for percentage of internet user at the district level, β_6 is percentage of poverty at the district level, and β_7 is the population density per square kilometer; v_k is a spatially unstructured random effect that is assumed to be normally distributed with mean zero and variance σ_v^2 and u_k is the spatially structured random effect that was modeled using a conditional autoregressive (CAR) prior $u_k \sim N(\bar{u}_{-k}, \sigma_u^2/n_k)$, where $-k$ denotes the neighbors of the k th district based on a simple adjacency matrix and n_k is the corresponding number of neighbors (7). WinBUGS software (version 1.4.3, MRC Biostatistics Unit, Cambridge, and Imperial College School of Medicine, London) was used to fit the Bayesian Poisson regression models. In the Markov chain Monte Carlo analysis, a 30,000 iteration “burn-in” was followed by 100,000 iteration sample collection. In every case, the Monte Carlo error was <5% of the overall standard deviation, indicating sufficient iterations of the model had been run after convergence. Model comparison was performed using the Deviance Information Criterion (DIC). Best fitted models were indicated by smaller DIC values (8).

The mean monthly number of pneumonia cases in children <5 years was 747.82. The mean monthly temperature, temperature variation, and rainfall were 30.97 °C, 3.63 °C, and 164.54 mm, respectively. Among the social factors, the mean percentages of education, internet use, poverty (per 100 population) and <5 years children density (per square kilometer) were 54.66%, 0.62%, 24.45%, and 117.72, respectively, at the district level (Supplementary Table S2, available in <http://weekly.chinacdc.cn>). The average monthly temperature was higher in the western region, while the monthly temperature variation was higher in most of the hilly areas located in the southern part of the country and two districts (Bhola and

Pirojpur) of the coastal region of Bangladesh. The distributions of higher monthly average rainfall were scattered in different regions. Inclusion of spatial autocorrelation in the model was important. The model which included both structured and unstructured random effects had the smallest DIC (665.47 and 13,773.00 for models with and without random effects, respectively) (Supplementary Table S3, available in <http://weekly.chinacdc.cn>). The highest RRs were observed in the southeastern part (Rangamati district) and southern part (Pirojpur district) of the country (Figure 1). Supplementary Table S4 (available in <http://weekly.chinacdc.cn>) shows the list of districts with the higher RR.

Figure 2 depicts the distribution of spatial random effects (structured heterogeneity) of pneumonia in Bangladesh. The districts with darker color (red color) had relatively high spatial variation. These districts with high spatial variation might have some unknown factors that may have had effects on the incidence of childhood pneumonia but that we did not consider in the models (e.g., incomplete measurement of variables, lack of geocoding, and generalization of geographic features).

Our study results suggested that a rise of 1 °C average monthly temperature and temperature




FIGURE 1. Posterior estimated Relative Risk of childhood pneumonia at the district level of Bangladesh from 2012 to 2016.




FIGURE 2. Spatial random effects of childhood pneumonia — Bangladesh, 2012–2016.

variation was associated with RR estimates of childhood pneumonia of 1.161 (95% CrI: 1.012–1.428) and 1.463 (95% CrI: 1.169–1.838), respectively. The density of children under five years in population was negatively associated with pneumonia (RR: 0.996, 95% CrI: 0.994–0.998) (Table 1).

TABLE 1. Crude and adjusted RR of different socioecological factors in Children Under Five-Years Old — Bangladesh, 2012–2016.

Variables	Crude RR (95% CrI)	Adjusted RR (95% CrI)
Temperature*	1.146 (0.929–1.432)	1.161 (1.013–1.429)
Temperature [†]	1.730 (1.694–1.763)	1.529 (1.503–1.555)
Temperature variability*	1.821 (1.376–2.491)	1.463 (1.170–1.839)
Temperature variability [†]	1.623 (1.596–1.649)	1.421 (1.395–1.447)
Rainfall*	1.000 (0.999–1.002)	1.000 (0.999–1.002)
Rainfall [†]	1.0007 (1.0006–1.0007)	1.0001 (1.0000–1.0002)
Population density*	0.995 (0.994–0.997)	0.996 (0.994–0.998)
Population density [†]	0.9943 (0.9942–0.9945)	0.9959 (0.9958–0.9961)
Education*	0.979 (0.956–1.004)	0.986 (0.969–1.005)
Education [†]	0.978 (0.977–0.979)	0.9853 (0.984–0.986)
Poverty*	1.008 (0.996–1.020)	1.000 (0.988–1.010)
Poverty [†]	1.009 (1.009–1.010)	1.002 (1.002–1.003)
Internet*	0.844 (0.661–1.070)	0.916 (0.743–1.126)
Internet [†]	0.874 (0.862–0.885)	0.929 (0.916–0.942)

Abbreviations: RR=relative risk; CrI=credible interval.

* with heterogeneity (u and v).

[†] without heterogeneity (u and v).

Additionally, no significant associations were found between childhood pneumonia and rainfall, education, internet use, or poverty since the corresponding 95% CrIs for the RR of each factor included 1.

DISCUSSION

In young children, the thermoregulation system is not yet matured and makes the children more vulnerable to temperature variation. This study describes the spatial pattern of childhood pneumonia and their socioecological factors in Bangladesh. Identifying the spatial variation of childhood pneumonia and important socioecological determinants can help target high-risk communities with evidence-based effective preventative measures.

Mapping of the spatially structured random effects indicated the spatial variation after controlling socioecological factors and spatial autocorrelation in the model. The Bayesian CAR model included unknown parameters as random effects, which incorporated the spatially correlated random effects (9). This approach can account for the residual variability resulting from spatial variation in effects that were not included in the models. The districts containing higher spatial random effects or variation may have some other risk factors remaining after adjustment of socioecological factors and spatial correlation.

This study was subject to some limitations. First, in this study, we used data from monthly reports of the DGHS. This represented the number of patients that attended different levels of health facilities in Bangladesh for pneumonia treatment. However, there might be some patients in the community who did not attend any health facilities and who received treatment from village doctors or spiritual healers, especially in the rural areas. Therefore, there was a chance of measurement and information biases. Second, the unit of analysis was at the group level rather than at the individual level, so the results may be prone to the ecological fallacy.

The findings of this study could help policymakers better understand that childhood pneumonia has a heterogeneous spatial pattern and that socioecological factors may play a significant role in describing this pattern.

Acknowledgements: The Management Information System, The Director General of Health Services, Ministry of Health, and Family Welfare, Bangladesh.

Funding: The Queensland University of Technology Postgraduate Research Award and Queensland University of Technology Higher Degree Research International Tuition Fee Sponsorship.

doi: 10.46234/ccdcw2021.161

* Corresponding author: Wenbiao Hu, w2.hu@qut.edu.au.

¹ School of Public Health and Social Work, Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Queensland, Australia; ² Infectious Diseases Division, International Centre for Diarrhoeal Disease Research, Bangladesh, Mohakhali, Dhaka, Bangladesh; ³ Shanghai Children's Medical Centre, School of Medicine, Shanghai Jiao Tong University, Shanghai, China; ⁴ School of Public Health, Institute of Environment and Population Health, Anhui Medical University, Hefei, Anhui, China; ⁵ Center for

Global Health, School of Public Health, Nanjing Medical University, Nanjing, Jiangsu, China; ⁶ Health System and Population Studies Division, International Centre for Diarrhoeal Disease Research, Bangladesh, Mohakhali, Dhaka, Bangladesh.

Submitted: May 12, 2021; Accepted: July 13, 2021

REFERENCES

1. DGHSIMCnewsletter2017 http://www.dghs.gov.bd/images/docs/IMCI/IMCI_12_2017.pdf. [2020-11-13].
2. World Health Organization. Revised WHO classification and treatment of childhood pneumonia at health facilities—evidence summaries. 2014. <https://apps.who.int/iris/rest/bitstreams/611120/retrieve>. [2021-1-22].
3. Bangladesh Bureau of Statistics. Population and household census. 2011. <http://www.bbs.gov.bd/site/page/47856ad0-7e1c-4aab-bd78-892733bc06eb/Population-and-Housing-Census>. [2021-2-18].
4. Bangladesh Bureau of Statistics, Statistics and Informatics Division, Ministry of Planning, Government of the People's Republic of Bangladesh. Bangladesh-household income and expenditure survey. 2016. expenditure survey. 2016. https://www.academia.edu/37874731/Preliminary_Report_on_Household_Income_and_Expenditure_Survey_2016_BANGLADESH_BUREAU_OF_STATISTICS-BBS_STATISTICS_AND_INFORMATICS_DIVISION_SID. [2020-9-17].
5. Gullón P, Varela C, Martínez EV, Gómez-Barroso D. Association between meteorological factors and hepatitis A in Spain 2010-2014. *Environ Int* 2017;102:230 – 5. <http://dx.doi.org/10.1016/j.envint.2017.03.008>.
6. Thomson MC, Connor SJ, D'Alessandro U, Rowlingson B, Diggle P, Cresswell M, et al. Predicting malaria infection in Gambian children from satellite data and bed net use surveys: the importance of spatial correlation in the interpretation of results. *Am J Trop Med Hyg* 1999;61(1):2 – 8. <http://dx.doi.org/10.4269/ajtmh.1999.61.2>.
7. Hu WB, Clements A, Williams G, Tong SL, Mengersen K. Spatial patterns and socioecological drivers of dengue fever transmission in Queensland, Australia. *Environ Health Perspect* 2012;120(2):260 – 6. <http://dx.doi.org/10.1289/ehp.1003270>.
8. Duncan EW, White NM, Mengersen K. Spatial smoothing in Bayesian models: a comparison of weights matrix specifications and their impact on inference. *Int J Health Geogr* 2017;16:47. <http://dx.doi.org/10.1186/s12942-017-0120-x>.
9. Best N, Richardson S, Thomson A. A comparison of bayesian spatial models for disease mapping. *Stat Methods Med Res* 2005;14(1):35 – 59. <http://dx.doi.org/10.1191/0962280205sm388oa>.

Supplementary Material

Statistical models

As a consequence, a total of 6 models were developed. Model I included only ecological measures (temperature, temperature variation, and rainfall) as explanatory variables; Model II included only social factors (education, internet use, population density, and poverty) as covariates; Model III included both ecological and social factors as explanatory variables; Model IV incorporated spatially structured random effects with all socioecological covariates; Model V incorporated spatially unstructured random effects with all socioecological covariates; Model VI incorporated both structured and unstructured random effects with all socioecological covariates.

The expected log relative risk θ_k for these models thus represented as follows:

$$\theta_k = \alpha + (\text{Temp}_k) \beta_1 + (\text{Tempva}_k) \beta_2 + (\text{Rain}_k) \beta_3 \dots \dots \dots \text{Model I}$$

$$\theta_k = \alpha + (\text{Edu}_k) \beta_1 + (\text{Int}_k) \beta_2 + (\text{povi}_k) \beta_3 + (\text{pop}_k) \beta_4 \dots \dots \dots \text{Model II}$$

$$\theta_k = \alpha + (\text{Temp}_k) \beta_1 + (\text{Tempva}_k) \beta_2 + (\text{Rain}_k) \beta_3 + (\text{Edu}_k) \beta_4 + (\text{Int}_k) \beta_5 + (\text{povi}_k) \beta_6 + (\text{pop}_k) \beta_7 \dots \dots \dots \text{Model III}$$

$$\theta_k = \alpha + (\text{Temp}_k) \beta_1 + (\text{Tempva}_k) \beta_2 + (\text{Rain}_k) \beta_3 + (\text{Edu}_k) \beta_4 + (\text{Int}_k) \beta_5 + (\text{povi}_k) \beta_6 + (\text{pop}_k) \beta_7 + u_k \dots \dots \dots \text{Model IV}$$

$$\theta_k = \alpha + (\text{Temp}_k) \beta_1 + (\text{Tempva}_k) \beta_2 + (\text{Rain}_k) \beta_3 + (\text{Edu}_k) \beta_4 + (\text{Int}_k) \beta_5 + (\text{povi}_k) \beta_6 + (\text{pop}_k) \beta_7 + v_k \dots \dots \dots \text{Model V}$$

$$\theta_k = \alpha + (\text{Temp}_k) \beta_1 + (\text{Tempva}_k) \beta_2 + (\text{Rain}_k) \beta_3 + (\text{Edu}_k) \beta_4 + (\text{Int}_k) \beta_5 + (\text{povi}_k) \beta_6 + (\text{pop}_k) \beta_7 + u_k + v_k \dots \dots \dots \text{Model VI}$$

where α is a constant, β_1 is the coefficient for temperature, β_2 is the coefficient for temperature variation, β_3 is the coefficient for rainfall, β_4 is the coefficient for percentage of education at district level, β_5 is the coefficient for percentage of internet user at district level, β_6 is percentage of poverty at district level, and β_7 is the population density per square kilometer, v_k is a spatially unstructured random effect that is assumed to be normally distributed with mean zero and variance σ_v^2 and u_k is spatially structured random effect that was modeled using a conditional autoregressive (CAR) prior $u_k \sim N(\bar{u}_{-k}, \sigma_u^2/n_k)$, where $-k$ denotes the neighbors of the k th district based on a simple adjacency matrix and n_k is the corresponding number of neighbors.

SUPPLEMENTARY TABLE S1. Spearman correlation between pneumonia and socioecological covariates in Children Under Five-Years Old — Bangladesh, 2012–2016.

Variables	1	2	3	4	5	6	7	8
1 Pneumonia	–							
2 Temperature	0.094							
3 Temperature variation	0.161	0.235						
4 Rainfall	–0.019	0.268*	–0.146					
5 Education	0.008	–0.011	–0.129	0.068				
6 Internet use	0.126	0.063	0.223	0.017	–0.066			
7 Population density	–0.148	–0.25*	–0.276*	0.028	0.075	0.162		
8 Poverty	0.069	–0.14	0.163	–0.077	0.095	–0.209	–0.336*	–

Note: – represent its pneumonia itself, there will be no number.

* $P < 0.05$.

SUPPLEMENTARY TABLE S2. Descriptive statistics of childhood pneumonia and different socioecological factors — Bangladesh, 2012–2016.

Variables	Mean \pm SD	Range
Pneumonia	747.82 \pm 245.32	355.53–1612.10
Temperature (°C)	30.97 \pm 0.48	29.50–31.98
Temperature variation (°C)	3.63 \pm 0.55	2.00–4.99
Rain (mm)	164.54 \pm 101.37	3.13–386.96
Education (%)	54.66 \pm 7.75	37.50–73.70
Internet use (%)	0.62 \pm 0.76	0.14–6.03
Poverty incidence (%)	27.45 \pm 15.31	2.60–70.80
Under five years population density (per square km)	117.72 \pm 87.05	10.81–656.54

SUPPLEMENTARY TABLE S3. Model comparison for relative risk of monthly childhood pneumonia, underlying socioecological factors, and different random effects — Bangladesh, 2012–2016.

Model	Random effect	Deviance Information Criterion (DIC)	Effective number of parameters (pD)
Model I	No	19954.20	30.146
Model II	No	17382.41	5.001
Model III	No	13773.00	11.134
Model IV	<i>u</i>	665.98	63.940
Model V	<i>v</i>	665.67	63.812
Model VI	<i>u and v</i>	665.47	63.719

SUPPLEMENTARY TABLE S4. List of high-risk districts of Bangladesh for childhood pneumonia from 2012 to 2016.

Name of the district	Relative Risk (95% Credible interval)	Location
Rangamati	5.97 (5.63–6.31)	South-eastern
Pirojpur	4.71 (4.48–4.93)	South-western
Jhalkathi	4.38 (4.09–4.66)	South-western
Jaipurh	3.95 (3.70–4.19)	North-eastern
Bandarban	3.77 (3.48–4.07)	South-eastern
Meherpur	3.50 (3.23–3.78)	South-western
Rajbari	3.31(3.11–3.50)	Central
Khagrachari	3.25 (3.02–3.49)	South-eastern
Panchagarh	2.96 (2.78–3.14)	Northern