

Supplementary Material

The rationale of using PM_{2.5}, O₃, NO₂, and SO₂ to construct the air quality health index (AQHI):

Similar to previous studies in Guangzhou Province (1) and New York City (2), the air pollutants in the AQHI included PM_{2.5}, O₃, NO₂, and SO₂. In Hong Kong, China, the developed AQHI were based on the health effects of PM₁₀, O₃, SO₂ (3), and NO₂. While in Canada, the established AQHI were based on the health effects of PM_{2.5}, O₃, and NO₂ (4). The inclusion of O₃ and NO₂ is not controversial. Considering the collinearity of PM_{2.5} and PM₁₀, and the health hazard differences between them, PM_{2.5} was chosen in our calculations. SO₂ was included by considering the pollution types in different regions of China. Several studies revealed that SO₂ was strongly associated with total mortality risk (5–6). SO₂ pollution mainly comes from the combustion of sulfur-containing fossil fuels (7). Notably, China is a country with massive coal production and consumption (8). According to the Bulletin of China's Ecological Environment 2018, the yearly average concentration of SO₂ in the Fen-Wei Plains was 24 µg/m³ in 2018 (9). According to our data in 2013–2018 (Supplementary Table S1), more than 25% of daily average concentration of county was higher than 26 µg/m³. Therefore, the health risk of SO₂ pollution in China still needs to be considered.

SUPPLEMENTARY TABLE S1. Frequency distribution of air quality health index (AQHI) in 36 cities^a of China, 2013–2018.

City	Level	Frequency	Distribution
Xiamen	[1,3]	1,381	68.64%
	(3,6]	609	30.27%
	(6,10]	22	1.09%
	(10,10+]	0	0%
Shenzhen	(1,3]	1,345	66.68%
	(3,6]	657	32.57%
	(6,10]	15	0.74%
	(10,10+]	0	0%
Dalian	(1,3]	959	47.62%
	(3,6]	968	48.06%
	(6,10]	84	4.17%
	(10,10+]	3	0.15%
Qingdao	(1,3]	705	34.88%
	(3,6]	1,183	58.54%
	(6,10]	121	5.59%
	(10,10+]	12	0.59%
Ningbo	(1,3]	748	37.12%
	(3,6]	1,172	58.16%
	(6,10]	92	4.57%
	(10,10+]	3	0.15%
Beijing	(1,3]	628	31.99%
	(3,6]	1,076	54.81%
	(6,10]	227	11.56%
	(10,10+]	32	1.63%
Shanghai	(1,3]	539	26.74%
	(3,6]	1,321	65.53%
	(6,10]	148	7.34%
	(10,10+]	8	0.40%

Continued

City	Level	Frequency	Distribution
Tianjin	(1,3]	398	19.77%
	(3,6]	1,234	61.30%
	(6,10]	317	15.75%
	(10,10+]	64	3.18%
Chongqing	(1,3]	704	34.90%
	(3,6]	1,273	63.11%
	(6,10]	40	1.98%
	(10,10+]	0	0%
Hefei	(1,3]	720	35.80%
	(3,6]	1,206	59.97%
	(6,10]	85	4.23%
	(10,10+]	0	0%
Fuzhou	(1,3]	1,502	74.58%
	(3,6]	508	25.22%
	(6,10]	4	0.20%
	(10,10+]	0	0%
Lanzhou	(1,3]	484	24.02%
	(3,6]	1,377	68.34%
	(6,10]	153	7.59%
	(10,10+]	1	0.05%
Guangzhou	(1,3]	798	39.62%
	(3,6]	1,086	53.92%
	(6,10]	125	6.21%
	(10,10+]	5	0.25%
Guilin	(1,3]	1,121	66.33%
	(3,6]	542	32.07%
	(6,10]	25	1.48%
	(10,10+]	2	0.12%
Guiyang	(1,3]	1375	68.24%
	(3,6]	603	29.93%
	(6,10]	37	1.84%
	(10,10+]	0	0%
Haikou	(1,3]	1,948	96.72%
	(3,6]	66	3.28%
	(6,10]	0	0%
	(10,10+]	0	0%
Shijiazhuang	(1,3]	250	12.39%
	(3,6]	1,113	55.18%
	(6,10]	463	22.95%
	(10,10+]	191	9.47%
Zhengzhou	(1,3]	1,345	66.68%
	(3,6]	657	32.57%
	(6,10]	15	0.74%
	(10,10+]	0	0%

Continued

City	Level	Frequency	Distribution
Harbin	(1,3]	789	39.12%
	(3,6]	882	43.73%
	(6,10]	264	13.09%
	(10,10+]	82	4.07%
Wuhan	(1,3]	493	24.52%
	(3,6]	1,198	59.57%
	(6,10]	293	14.57%
	(10,10+]	27	1.34%
Changsha	(1,3]	818	40.62%
	(3,6]	1,050	52.14%
	(6,10]	146	7.25%
	(10,10+]	0	0%
Changchun	(1,3]	660	32.77%
	(3,6]	1,126	55.91%
	(6,10]	212	10.53%
	(10,10+]	16	0.79%
Nanjing	(1,3]	438	21.73%
	(3,6]	1,339	66.42%
	(6,10]	228	11.31%
	(10,10+]	11	0.55%
Nanchang	(1,3]	973	48.31%
	(3,6]	963	47.82%
	(6,10]	75	3.72%
	(10,10+]	3	0.15%
Shenyang	(1,3]	402	20.00%
	(3,6]	1,181	58.76%
	(6,10]	324	16.12%
	(10,10+]	103	5.12%
Hohhot	(1,3]	538	26.74%
	(3,6]	1,229	61.08%
	(6,10]	237	11.78%
	(10,10+]	8	0.40%
Yinchuan	(1,3]	530	26.38%
	(3,6]	1,108	55.15%
	(6,10]	315	15.68%
	(10,10+]	56	2.79%
Xining	(1,3]	553	27.43%
	(3,6]	1,404	69.64%
	(6,10]	59	2.93%
	(10,10+]	0	0%
Jinan	(1,3]	133	6.58%
	(3,6]	1,385	68.50%
	(6,10]	421	20.82%
	(10,10+]	83	4.10%

Continued

City	Level	Frequency	Distribution
Taiyuan	(1,3]	320	15.90%
	(3,6]	1,198	59.54%
	(6,10]	399	19.83%
	(10,10+]	95	4.72%
Xi'an	(1,3]	365	18.13%
	(3,6]	1,344	66.77%
	(6,10]	264	13.11%
	(10,10+]	40	1.99%
Chengdu	(1,3]	387	19.22%
	(3,6]	1,422	70.61%
	(6,10]	198	9.83%
	(10,10+]	7	0.35%
Lhasa	(1,3]	1,792	88.98%
	(3,6]	222	11.02%
	(6,10]	0	0%
	(10,10+]	0	0%
Urumchi	(1,3]	530	26.36%
	(3,6]	1,161	57.73%
	(6,10]	292	14.52%
	(10,10+]	28	1.39%
Kunming	(1,3]	1,255	62.31%
	(3,6]	755	37.49%
	(6,10]	4	0.20%
	(10,10+]	0	0%
Hangzhou	(1,3]	709	35.19%
	(3,6]	1,228	60.94%
	(6,10]	76	3.77%
	(10,10+]	2	0.10%

Note: 36 cities refer to all municipalities directly under the central government, cities specifically designated in the state plan, and provincial-level capitals.

REFERENCES

- Li X, Xiao JP, Lin HL, Liu T, Qian ZM, Zeng WL, et al. The construction and validity analysis of AQHI based on mortality risk: A case study in Guangzhou, China. *Environ Pollut* 2017;220:487 – 94. <http://dx.doi.org/10.1016/j.envpol.2016.09.091>.
- Perlmutter LD, Cromar KR. Comparing associations of respiratory risk for the EPA Air Quality Index and health-based air quality indices. *Atmos Environ* 2019;202:1 – 7. <http://dx.doi.org/10.1016/j.atmosenv.2019.01.011>.
- Wong TW, Tam WWS, Yu ITS, Lau AKH, Pang SW, Wong AHS. Developing a risk-based air quality health index. *Atmos Environ* 2013;76:52 – 8. <http://dx.doi.org/10.1016/j.atmosenv.2012.06.071>.
- Stieb DM, Burnett RT, Smith-Doiron M, Brion O, Shin HH, Economou V. A new multipollutant, no-threshold air quality health index based on short-term associations observed in daily Time-Series analyses. *J Air Waste Manag Assoc* 2008;58(3):435 – 50. <http://dx.doi.org/10.3155/1047-3289.58.3.435>.
- Hoek G, Brunekreef B, Verhoeff A, van Wijnen J, Fischer P. Daily mortality and air pollution in the Netherlands. *J Air Waste Manag Assoc* 2000;50(8):1380 – 9. <http://dx.doi.org/10.1080/10473289.2000.10464182>.
- World Health Organization. Air quality guidelines-global update 2005. *Public Health & Environment* 2014. https://www.who.int/phe/health_topics/outdoorair/outdoorair_aqg/en/. [2021-01-04].
- Tang XY, Zhang YH, Shao M. Atmospheric environmental chemistry. 2nd ed. Beijing: China Higher Education Press. 2006;p.269 – 274.
- Wang Q, Song XX, Liu Y. China's coal consumption in a globalizing world: insights from multi-regional input-output and structural decomposition analysis. *Sci Total Environ*. 2020;711:134790. <http://dx.doi.org/10.1016/j.scitotenv.2019.134790>.
- Ministry of Ecology and Environment of the People's Republic of China. Bulletin on the State of the Ecological Environment in China. http://www.mee.gov.cn/xxgk2018/xxgk/xxgk15/201912/t20191231_754139.html. (In Chinese).