

Vital Surveillances

Injury Mortality of Children and Adolescents Aged 0–19 Years — China, 2010–2021

Jingtao Zhou¹; Min Zhao¹; Hao Huang¹; David C. Schwebel²; Peishan Ning¹; Zhenzhen Rao¹;
Peixia Cheng³; Li Li^{1,4}; Guoqing Hu^{1,4,5}

ABSTRACT

Introduction: To examine the recent trends in child injury mortality in China.

Methods: Injury mortality data of 2010–2021 for children and adolescents aged 0–19 years were from the China Health Statistics Yearbook. Injury mortality disparities across urban vs. rural locations, gender, and age groups were scrutinized. Annual percent change (APC), average annual percent change (AAPC), and their 95% confidence intervals (95% CI) were estimated using Joinpoint regression models.

Results: The age-standardized injury mortality significantly dropped from 21.87 to 9.41 per 100,000 population among children and youth aged 0–19 years during 2010–2021, with an AAPC of –6.7% (95% CI: –8.2%, –5.2%). The urban-rural disparity and gender gap in injury mortality reduced gradually. In 2021, drowning and road traffic crashes were the top two causes of child injury deaths, explaining 31.1% and 27.9% of total injury deaths, respectively. Suffocation accounted for 62.3% of injury deaths among infants younger than a year. Alarming, the suicide mortality rate rose from 2.16 to 3.42 per 100,000 population between 2010 and 2021 among teenagers aged 15–19 years. Subgroup analyses yielded similar results.

Conclusions: During 2010–2021, the injury mortality decreased significantly among Chinese children and adolescents, and the responding urban-rural disparities narrowed.

Injuries pose a significant global challenge to the health and safety of children and adolescents (1). According to estimates by the Global Burden of Disease (GBD) study group, injuries caused 595,621 deaths and 233,114,563 incident cases worldwide among individuals aged 0–19 years in 2019. Of these, 6.5% of the deaths and 9.3% of the incident cases

occurred in China (2).

The United Nations (UN) has outlined several Sustainable Development Goals (SDGs) pertinent to preventing injuries amongst children and adolescents (3). In the same vein, the Chinese central government promulgated a series of developmental outlines for children, and the current outline (2021–2030) sets an ambitious goal of a 20% reduction in child injury mortality by 2030 compared to 2020 (4).

To effectively monitor the progress towards the specified targets in China, it is critical to regularly analyze nationwide data to discern patterns in child and adolescent mortality due to injuries. Several studies have documented trends in this field until 2020. Research by Luo et al. (5) analyzed the temporal progression of mortality due to injuries in children and adolescents aged 1–24 years, utilizing the Chinese Cause of Death Surveillance dataset spanning from 2010 to 2020. Zheng et al. (6) reported a notable reduction in mortality from drowning and road traffic crashes in children and adolescents aged 5–19 years between 2008 and 2019. Elevated injury mortality were reported occurring among boys, children, and early adolescents aged 5–14, and individuals residing in the western and rural areas. Yao et al. explored disparities in injury-induced mortality among children by gender and area (urban vs. rural) in Sichuan Province (7), but their findings were not nationally representative.

This study scrutinized nationally representative data to assess trends in overall injury mortality rates among Chinese children and adolescents (aged 0–19 years) from 2010 to 2021. In addition, we evaluated subgroup mortality rates according to demographic factors including area (rural vs. urban), sex, and age group.

METHODS

This study sourced annual data on child and adolescent injury mortality from the Chinese Health

Statistical Yearbook (2010–2021), which provides age-specific mortality data for 11 categories of unintentional and intentional injuries (8): motor vehicle crashes, non-motor vehicle crashes, poisoning, falls, fire/burn injuries, drowning, mechanical suffocation, falling object injuries, electric shocks, homicide, and suicide. To estimate the year-end population for each year from 2010 to 2021, we employed linear interpolation (9) using China's census data of 2010 and 2020. The population data from 2020 was then utilized as the standard population for calculating age-standardized injury mortality.

Linear graphs were utilized to display variations in both overall and subgroup injury mortality, separated by area, gender, and age group for Chinese children and adolescents aged 0 to 19 years. Furthermore, stacked area charts were constructed to show the cause spectrum of child and adolescent injury mortality from 2010 to 2021, segmented by area and age group.

Significant injury mortality changes throughout the study time period were quantified using average annual percent change (AAPC) and annual percent change (APC), alongside their corresponding 95% confidence

intervals (95% CIs), which were estimated via Joinpoint regression models. We used the Joinpoint Regression Program (Version 4.9.1.0, National Cancer Institute, Calverton, USA) to perform statistical analysis.

RESULTS

Overall Injury Mortality and Subgroup Mortality by Area and Age Group

Between 2010 and 2021, the overall age-standardized injury mortality for children and adolescents in China decreased from 21.87 to 9.41 per 100,000 population (AAPC=−6.7%, 95% CI: −8.2%, −5.2%). The age-standardized injury mortality for urban children was continuously lower than and declined more slowly than that for rural children (urban–rural mortality ratio: 0.53 to 0.70; AAPC: −4.6% *vs.* −6.9%) (Figure 1A and Table 1). Injury mortality declined significantly during 2010–2021 in four age groups (under 1 year: AAPC=−7.9%, 1–4 years: AAPC=−11.6%, 5–9 years: AAPC=−7.5%,




FIGURE 1. Age-standardized and age-specific mortality rates for injuries among children and adolescents in various regions of China, from 2010 to 2021. (A) Age-standardized injury mortality rates in individuals aged 0–19 segmented by area. (B) Mortality caused by injuries in children under 1 year, differentiated by area. (C) Injury-caused mortality in children aged 1–4, broken down by area. (D) Injury-associated mortality in children aged 5–9, classified by area. (E) Mortality due to injuries in adolescents aged 10–14, outlined by area. (F) Injury mortality trends in adolescents aged 15–19, divided by area.

TABLE 1. APC and AAPC in injury mortality among Chinese children and adolescents from 2010 to 2021.

Age group	Area	Block 1		Block 2		Block 3		AAPC (95% CI)
		Time period	APC (95% CI)	Time period	APC (95% CI)	Time period	APC (95% CI)	
All ages	Total	2010–2021	–6.7 (–8.2, –5.2)*					–6.7 (–8.2, –5.2)*
	Urban	2010–2021	–4.6 (–6.5, –2.6)*					–4.6 (–6.5, –2.6)*
	Rural	2010–2021	–6.9 (–8.4, –5.5)*					–6.9 (–8.4, –5.5)*
0–1 years	Total	2010–2021	–7.9 (–9.7, –6.1)*					–7.9 (–9.7, –6.1)*
	Urban	2010–2021	–7.2 (–9.7, –4.6)*					–7.2 (–9.7, –4.6)*
	Rural	2010–2012	15.9 (–1.8, 36.8)	2012–2015	–14.1 (–27.3, 1.4)	2015–2021	–7.0 (–9.5, –4.3)*	–5.3 (–9.0, –1.4)*
1–4 years	Total	2010–2021	–11.6 (–14.1, –9.0)*					–11.6 (–14.1, –9.0)*
	Urban	2010–2021	–8.8 (–11.1, –6.4)*					–8.8 (–11.1, –6.4)*
	Rural	2010–2021	–11.7 (–14.6, –8.6)*					–11.7 (–14.6, –8.6)*
5–9 years	Total	2010–2021	–7.5 (–9.6, –5.3)*					–7.5 (–9.6, –5.3)*
	Urban	2010–2013	–9.6 (–9.4, 32.6)	2013–2021	–8.6 (–12.3, –4.7)*			–3.9 (–8.6, 1.0)
	Rural	2010–2014	–8.0 (–10.4, –5.6)*					–8.0 (–10.4, –5.6)*
10–14 years	Total	2010–2013	18.1 (6.7, 30.6)*	2013–2021	–5.4 (–7.5, –3.3)*			0.5 (–2.1, 3.2)
	Urban	2010–2013	17.0 (2.1, 34)*	2013–2021	–4.7 (–7.5, –1.9)*			0.7 (–2.8, 4.4)
	Rural	2010–2013	20.1 (8.1, 33.4)*	2013–2021	–5.3 (–7.4, –3.1)*			1.1 (–1.7, 3.9)
15–19 years	Total	2010–2021	–3.6 (–5.9, –1.3)*					–3.6 (–5.9, –1.3)*
	Urban	2010–2021	–2.1 (–4.8, 0.7)					–2.1 (–4.8, 0.7)
	Rural	2010–2014	4.6 (–7.0, 17.7)	2014–2018	–11.1 (–26.2, –7.1)	2018–2021	1.5 (–15.7, 22.3)	–2.2 (–8.5, 4.6)

Abbreviation: APC=annual percent change; AAPC=average annual percent change; CI=confidence interval.

* $P < 0.05$.

15–19 years: AAPC=–3.6%), while significant decreases only occurred between 2013 and 2021 for the age group of 10–14 years (Figure 1B–F and Table 1).

Subgroup Injury Mortality by Age Group, Sex, and Area

Between 2010 and 2021, age-standardized injury mortality decreased from 28.52 to 11.24 per 100,000 population for boys, and decreased from 14.34 to 7.30 per 100,000 population for girls. The age-standardized injury mortality for boys was continuously higher than and declined faster than that for girls in both urban and rural areas (Figure 2A and Table 2).

Subgroup analyses showed significant injury mortality decreases in three younger age groups for both sexes in both urban and rural areas (under 1 year, aged 1–4 years and aged 5–9 years, with AAPCs ranging from –12.8% to –4.3%), with exceptions for rural girls under 1 year and for urban boys aged 5–9 years old. Strikingly, overall and subgroup injury mortality by area and sex did not change significantly among early adolescents aged 10–14 years old. Significant reductions appeared only in urban boys (AAPC=–3.5%, 95% CI: –6.0%, –1.0%) and rural

girls (AAPC=–2.1%, 95% CI: –4.0%, –0.1%) for the age group of 15–19 years (Figure 2B–D, Table 2, and Supplementary Tables S1–S3, available at <https://weekly.chinacdc.cn/>).

Cause Spectrum of Injury Mortality by Age Group and Area

Supplementary Figure S1 (available at <https://weekly.chinacdc.cn/>) depicts the distribution of the top six causes of injury-related deaths, segmented by age group and geographical area. Drowning and road traffic crashes were the predominant causes of injury-related mortality for the four older age groups. Among infants under the age of 1 year, suffocation emerged as the primary cause, accounting for 56.0%–76.7% of injury deaths during 2010–2021. Suicide has come to light as a significant cause of injury-related deaths among adolescents aged 10–19 years.

During 2010–2021, significant injury mortality decreases were detected for suffocation among children under 1 year (urban: AAPC=–7.7%, rural: AAPC=–6.8%). Drowning mortality decreased significantly among children and adolescents aged 0–19 years (urban: AAPC=–8.0%, rural: AAPC=–9.5%). The




FIGURE 2. Overall age-standardized and age-specific mortality rates due to injury among Chinese children and adolescents, categorized by sex and area, spanning the years 2010–2021. (A) Age-standardized injury mortality in children and adolescents aged 0–19, separated by sex and area. (B) Injury mortality in infants under one year of age, reported by sex and area. (C) Injury mortality in toddlers aged 1–4, delineated by sex and area. (D) Injury mortality in young children aged 5–9, categorized by sex and area. (E) Injury mortality in early adolescents aged 10–14, divided by sex and area. (F) Injury mortality in late adolescents aged 15–19, sorted by sex and area.

overall and age-specific road traffic injury mortality reduced substantially since 2014 (e.g., 0–19 years, urban area: APC=−10.6%, rural area: APC=−10.2%). In contrast, notable suicide mortality increases were observed in the two oldest age groups for both urban and rural areas (with an AAPC changing between 4.7% and 11.5%) (Supplementary Figure S1 and Supplementary Table S4, available at <https://weekly.chinacdc.cn/>).

DISCUSSION

This study, using nationally representative data, analyzed current trends in injury mortality among Chinese children aged 0–19 years. The research produced four critical findings: First, the overall age-standardized injury mortality saw a decrease of 57% between 2010 and 2021. Second, the mortality rate due to injuries was comparatively higher in rural areas and among boys, with a faster decline rate than in urban areas and among girls. Third, the predominant cause of injury mortality differed by age groups, where

unintentional suffocation was the leading cause of death for infants under one year and drowning was the main cause for the four older age groups. Lastly, suicide emerged as a leading cause of injury mortality among adolescents aged 10–19 years.

The significant decrease in injury mortality rates in Chinese children and adolescents over the past ten years are likely to be attributed to two main factors. First, the reduction may symbolize the enormous efforts made by the Chinese government regarding injury prevention. As an instance, the implementation of changes to traffic laws — such as making drunk driving a criminal offense — has decreased traffic crash risks (10). Furthermore, the Chinese administration issued a variety of injury prevention guidelines and commenced education programs in schools across the country. These initiatives probably have a positive impact on child injury prevention.

Second, the notable decline in mortality due to injuries might be correlated with the swift urbanization throughout the nation. The percentage of Chinese residents residing in rural areas dropped from 48.17%

TABLE 2. Mortality due to injuries and AAPC among Chinese children and adolescents, categorized by sex, from 2010–2021.

Age group	Area	Boys			Girls		
		Mortality in 2010	Mortality in 2021	AAPC (95% CI)	Mortality in 2010	Mortality in 2021	AAPC (95% CI)
All ages	Total	28.52	11.24	−7.4 (−9.0, −5.7)*	14.34	7.30	−5.6 (−6.9, −4.2)*
	Urban	18.62	9.61	−5.3 (−7.4, −3.1)*	9.97	6.61	−3.4 (−5.1, −1.7)*
	Rural	36.14	14.53	−7.4 (−9.1, −5.8)*	17.75	8.56	−6.0 (−7.4, −4.7)*
Under 1 year	Total	41.05	16.82	−8.0 (−9.4, −6.6)*	27.15	16.96	−7.7 (−10.5, −4.8)*
	Urban	36.21	15.19	−7.0 (−9.7, −4.2)*	28.74	14.87	−7.5 (−10.5, −4.5)*
	Rural	44.15	19.62	−7.5 (−8.8, −6.2)*	26.10	20.58	−3.5 (−13.0, 7.0)
1–4 years	Total	66.00	10.41	−11.9 (−14.7, −9.0)*	38.40	7.79	−11.0 (−13.1, −8.8)*
	Urban	33.42	8.51	−9.0 (−11.5, −6.4)*	21.93	6.79	−8.4 (−10.4, −6.2)*
	Rural	89.24	14.28	−11.9 (−15.2, −8.5)*	50.31	9.82	−12.8 (−15.9, −9.7)*
5–9 years	Total	16.45	6.65	−8.3 (−10.3, −6.3)*	7.53	4.05	−5.9 (−8.7, −2.9)*
	Urban	11.29	5.80	−4.5 (−9.3, 0.6)	6.00	3.41	−4.3 (−7.6, −0.8)*
	Rural	20.42	8.14	−9.0 (−11.0, −6.9)*	8.71	5.16	−6.1 (−9.2, −2.8)*
10–14 years	Total	12.49	11.08	−0.8 (−3.6, 2.1)	4.67	6.75	3.0 (−2.0, 8.2)
	Urban	10.02	8.99	−0.6 (−3.5, 2.3)	3.85	6.03	2.3 (−1.3, 6.2)
	Rural	14.45	14.21	−0.8 (−4.9, 3.5)	5.32	7.83	3.3 (−1.2, 8.0)
15–19 years	Total	26.26	17.03	−4.8 (−7.1, −2.5)*	10.22	9.96	−0.9 (−3.8, 2.1)
	Urban	21.54	15.19	−3.5 (−6.0, −1.0)*	8.17	9.79	1.1 (−2.6, 5.0)
	Rural	31.67	22.28	−3.6 (−7.2, 0.2)	12.66	10.52	−2.1 (−4.0, −0.1)*

Abbreviation: AAPC=average annual percent change; CI=confidence interval.

* $P < 0.05$.

in 2011 to 35.28% in 2021 (11). This could possibly lead to lower exposure to specific injury risks and hazards, such as drowning, for children.

Our results corroborate earlier findings that injury-related death rates are significantly elevated among children in rural regions and boys, compared to their urban and female counterparts (5). The heightened injury mortality rates among young people in China's rural areas are linked to insufficient adult supervision of children who are left behind (12), as well as a lack of adequate public facilities including readily accessible medical centers. The higher injury fatality rates among boys could be attributed to their increased physical activity, impulsivity, and heightened energy levels, predisposing them to engage in high-risk behaviors (5).

The substantial decrease in injury mortality rates in rural areas between 2010 and 2021 is promising. It may be attributable to accelerated socio-economic advancement experienced in rural China over the past decade. The National Precision Poverty Alleviation Project has seen considerable progress in strengthening rural economic growth and distribution of healthcare resources. From 2010 to 2021, there was an impressive 115.8% rise in the number of health technicians per

thousand population in rural sectors, greatly surpassing the 32.9% increase observed in urban sectors (8). Furthermore, the count of children left behind in rural primary and middle schools declined from 15.51 million in 2017 to 12.00 million in 2021 (11).

Our findings indicate that the reasons for injury-related mortalities vary among the five age groups, reflecting different levels of exposure to certain hazards during various stages of child development. Infants under one-year-old, for instance, are more prone to unintentional suffocation due to underdeveloped respiratory systems, making them susceptible to potentially fatal obstructions while eating, playing, or sleeping (13). Older children, on the other hand, are more likely to engage in dangerous activities when left unsupervised, especially near roads or bodies of water (1). During adolescence, the rapid maturation of the brain and hormonal changes coupled with increased exposure to internet and social media, may exacerbate interpersonal stress and emotional instability (14). Such factors could potentially contribute to a rise in suicide rates among adolescents (14). The escalating trend of adolescent suicides that we have observed in China mirrors a global pattern, although the reported

rates in China remain lower than in many other countries (15).

Our findings have two policy implications. First, the substantial injury mortality reductions suggest that government goals to reduce child and adolescent injury mortality between 2011 and 2020 have been successful in China. Reduced injury mortality gaps between urban and rural areas and between boys and girls indicate slight improvement in reducing disparities and achieving injury mortality equity across population subgroups.

Second, systematic and intensified prevention efforts should continue according to prevention priorities listed by the Child Development Program of China (2021–2030). Priority should be identification of a government department responsible for injury prevention programs in China to lead implementation of proven prevention programs nationwide like the recent adoption of national child safety seat law.

This study has several limitations. First, due to the absence of data on non-fatal injuries, the results for injury morbidity over the past decade may differ. Second, because data were lacking, this research did not study relevant influencing factors. Conducting research to identify associated risk factors is necessary to fully interpret epidemiological data trends, quantify causal relations between influencing factors and injury mortality, and develop prevention programs.

CONCLUSIONS

From 2010 to 2021, there was a significant decline in child injury mortality in China. Moreover, injury mortality disparities across area (urban vs. rural) and gender (boys vs. girls) diminished during this period. The dramatic injury mortality decrease likely echoes the impact of governmental interventions. Considering the notably high child injury mortality, however, comprehensive and intensified efforts are encouraged to meet the objectives set forth by the Child Development Program of China (2021–2030).

Conflicts of interest: No conflicts of interest.

Funding: Supported by the National Natural Science Foundation of China (grant numbers 82073672, 82273743, and 82204165).

doi: 10.46234/ccdcw2024.057

* Corresponding authors: Li Li, lili1009@csu.edu.cn; Guoqing Hu, huguqing009@gmail.com.

¹ Department of Epidemiology and Health Statistics, Hunan Provincial Key Laboratory of Clinical Epidemiology, Xiangya School of Public

Health, Central South University, Changsha City, Hunan Province, China; ² Department of Psychology, University of Alabama at Birmingham, Birmingham, AL, USA; ³ Department of Child, Adolescent and Women's Health, School of Public Health, Capital Medical University, Beijing, China; ⁴ National Clinical Research Center for Geriatric Disorders, Xiangya Hospital, Central South University, Changsha City, Hunan Province, China.

Submitted: November 20, 2023; Accepted: January 30, 2024

REFERENCES

1. World Health Organization. World report on child injury prevention. Geneva: World Health Organization. 2008.
2. Institute for Health Metrics and Evaluation. GBD Results. 2019. <https://vizhub.healthdata.org/gbd-results/>. [2023-7-26].
3. United Nations Development Programme. Sustainable development goals. <https://www.undp.org/sustainable-development-goals/good-health>. [2023-7-26].
4. National Working Committee on Children and Women under State Council. China National Program for Child Development (2021–2030). 2021. <https://www.nwccw.gov.cn/2021/09/27/99338976.html>. [2023-7-26]. (In Chinese).
5. Luo XB, Zhang L, Wang WJ, Yang JL, Chang Y. Epidemiological characteristics of injury deaths among children and adolescents in China, 2010–2020. *Chin J Sch Health* 2023;44(8):1247 – 51. <https://doi.org/10.16835/j.cnki.1000-9817.2023.08.029>.
6. Zheng JH, Feng GS, Wu XF, Yu SC, Wang QQ. Mortality of drowning and road traffic injury among children aged 5–14 in China from 2008 to 2019. *Chin J Prev Med* 2022;56(9):1244 – 50. <https://doi.org/10.3760/cma.j.cn112150-20220212-00130>.
7. Yao MH, Wu GH, Zhao ZL, Luo M, Zhang JY. Unintentional injury mortality among children under age five in urban and rural areas in the Sichuan province of west China, 2009–2017. *Sci Rep* 2019;9(1):2963. <https://doi.org/10.1038/s41598-019-38936-6>.
8. National Health Commission of the People's Republic of China. Chinese Health Statistical Yearbook 2022. Beijing: Peking Union Medical College Press. 2022. <http://www.nhc.gov.cn/mohwsbwstjxxzx/tjtnj/202305/6ef68aac6bd14c1eb9375e01a0faa1fb.shtml>. (In Chinese).
9. Wang Y, Zhang X, Lu H, Matthews KA, Greenlund KJ. Intercensal and postcensal estimation of population size for small geographic areas in the United States. *Int J Popul Data Sci* 2020;5(1):1160. <https://doi.org/10.23889/ijpds.v5i1.1160>.
10. Xu XH, Dong H, Li L, Yang Z, Lin GZ, Ou CQ. Time-varying effect of drunk driving regulations on road traffic mortality in Guangzhou, China: an interrupted time-series analysis. *BMC Public Health* 2021;21(1):1885. <https://doi.org/10.1186/s12889-021-11958-4>.
11. National Bureau of Statistics of China. China statistical yearbook 2022. Beijing: China Statistics Press. 2022. <http://www.stats.gov.cn/sj/ndsj/2022/indexch.htm>.
12. Hu HW, Gao JM, Jiang HC, Xing PN. A comparative study of unintentional injuries among schooling left-behind, migrant and residential children in China. *Int J Equity Health* 2018;17(1):47. <https://doi.org/10.1186/s12939-018-0767-3>.
13. Yu X, Miao L, Zhu J, Liang J, Dai L, Li XH, et al. Social and environmental risk factors for unintentional suffocation among infants in China: a descriptive analysis. *BMC Pediatr* 2021;21(1):465. <https://doi.org/10.1186/s12887-021-02925-4>.
14. Clayton MG, Pollak OH, Prinstein MJ. Why suicide? Suicide propinquity and adolescent risk for suicidal thoughts and behaviors. *Clin Child Fam Psychol Rev* 2023;26(4):904 – 18. <https://doi.org/10.1007/s10567-023-00456-1>.
15. Glenn CR, Kleiman EM, Kellerman J, Pollak O, Cha CB, Esposito EC, et al. Annual research review: A meta-analytic review of worldwide suicide rates in adolescents. *J Child Psychol Psychiatry* 2020;61(3):294 – 308. <https://doi.org/10.1111/jcpp.13106>.

SUPPLEMENTARY MATERIALS

SUPPLEMENTARY TABLE S1. APC and AAPC in injury-related mortality among Chinese boys from 2010 to 2021.

Age group	Area	Block 1		Block 2		AAPC (95% CI)
		Time period	APC (95% CI)	Time period	APC (95% CI)	
0–19 years	Total	2010–2021	–7.4 (–9.0, –5.7)*			–7.4 (–9.0, –5.7)*
	Urban area	2010–2021	–5.3 (–7.4, –3.1)*			–5.3 (–7.4, –3.1)*
	Rural area	2010–2021	–7.4 (–9.1, –5.8)*			–7.4 (–9.1, –5.8)*
under 1 year	Total	2010–2021	–8.0 (–9.4, –6.6)*			–8.0 (–9.4, –6.6)*
	Urban area	2010–2021	–7.0 (–9.7, –4.2)*			–7.0 (–9.7, –4.2)*
	Rural area	2010–2021	–7.5 (–8.8, –6.2)*			–7.5 (–8.8, –6.2)*
1–4 years	Total	2010–2021	–11.9 (–14.7, –9.0)*			–11.9 (–14.7, –9.0)*
	Urban area	2010–2021	–9.0 (–11.5, –6.4)*			–9.0 (–11.5, –6.4)*
	Rural area	2010–2021	–11.9 (–15.2, –8.5)*			–11.9 (–15.2, –8.5)*
5–9 years	Total	2010–2021	–8.3 (–10.3, –6.3)*			–8.3 (–10.3, –6.3)*
	Urban area	2010–2013	8.8 (–10.8, 32.6)	2013–2021	–9.1 (–12.9, –5.0)*	–4.5 (–9.3, 0.6)
	Rural area	2010–2021	–9.0 (–11.0, –6.9)*			–9.0 (–11.0, –6.9)*
10–14 years	Total	2010–2013	17.7 (5.5, 31.3)*	2013–2021	–6.9 (–9.1, –4.7)*	–0.8 (–3.6, 2.1)
	Urban area	2010–2013	17.0 (4.6, 30.8)*	2013–2021	–6.5 (–8.7, –4.2)*	–0.6 (–3.5, 2.3)
	Rural area	2010–2014	13.1 (0.9, 26.6)*	2014–2021	–7.9 (–12.2, –3.4)*	–0.8 (–4.9, 3.5)
15–19 years	Total	2010–2021	–4.8 (–7.1, –2.5)*			–4.8 (–7.1, –2.5)*
	Urban area	2010–2021	–3.5 (–6.0, –1.0)*			–3.5 (–6.0, –1.0)*
	Rural area	2010–2013	8.2 (–6.7, 25.5)	2013–2021	–7.6 (–10.6, –4.6)*	–3.6 (–7.2, 0.2)

Abbreviation: APC=annual percent change; AAPC=average annual percent change; CI=confidence interval.

* $P<0.05$.

SUPPLEMENTARY TABLE S2. APC and AAPC in injury-induced mortality among Chinese girls spanning 2010 through 2021.

Age group	Area	Block 1		Block 2		Block 3		AAPC (95% CI)
		Time period	APC (95% CI)	Time period	APC (95% CI)	Time period	APC (95% CI)	
0–19 years	Total	2010–2021	-5.6 (-6.9, -4.2)*					-5.6 (-6.9, -4.2)*
	Urban area	2010–2021	-3.4 (-5.1, -1.7)*					-3.4 (-5.1, -1.7)*
	Rural area	2010–2021	-6.0 (-7.4, -4.7)*					-6.0 (-7.4, -4.7)*
Under 1 year	Total	2010–2021	-7.7 (-10.5, -4.8)*					-7.7 (-10.5, -4.8)*
	Urban area	2010–2021	-7.5 (-10.5, -4.5)*					-7.5 (-10.5, -4.5)*
	Rural area	2010–2012	33.8 (-13.1, 105.9)	2012–2015	-17.4 (-46.3, 27.2)	2015–2021	-6.5 (-13.1, 0.6)	-3.5 (-13.0, 7.0)
1–4 years	Total	2010–2021	-11.0 (-13.1, -8.8)*					-11.0 (-13.1, -8.8)*
	Urban area	2010–2021	-8.4 (-10.4, -6.2)*					-8.4 (-10.4, -6.2)*
	Rural area	2010–2012	-28.9 (-42.8, -11.6)*	2012–2021	-8.8 (-10.6, -7.0)*			-12.8 (-15.9, -9.7)*
5–9 years	Total	2010–2021	-5.9 (-8.7, -2.9)*					-5.9 (-8.7, -2.9)*
	Urban area	2010–2021	-4.3 (-7.6, -0.8)*					-4.3 (-7.6, -0.8)*
	Rural area	2010–2021	-6.1 (-9.2, -2.8)*					-6.1 (-9.2, -2.8)*
10–14 years	Total	2010–2013	18.4 (-2.0, 43.1)	2013–2021	-2.3 (-6.3, 1.8)			3.0 (-2.0, 8.2)
	Urban area	2010–2021	2.3 (-1.3, 6.2)					2.3 (-1.3, 6.2)
	Rural area	2010–2013	21.0 (2.2, 43.3)*	2013–2021	-2.7 (-6.2, 1.0)			3.3 (-1.2, 8.0)
15–19 years	Total	2010–2021	-0.9 (-3.8, 2.1)					-0.9 (-3.8, 2.1)
	Urban area	2010–2021	1.1 (-2.6, 5.0)					1.1 (-2.6, 5.0)
	Rural area	2010–2021	-2.1 (-4.0, -0.1)*					-2.1 (-4.0, -0.1)*

Abbreviation: APC=annual percent change; AAPC=average annual percent change; CI=confidence interval.

* $P < 0.05$.

SUPPLEMENTARY TABLE S3. AAPC in mortality rates due to injuries among Chinese children and adolescents, segregated by age group and region, from 2010 to 2021.

Age group	Area	Overall injury mortality			Injury mortality for boys			Injury mortality for girls		
		2010	2021	AAPC (95% CI)	2010	2021	AAPC (95% CI)	2010	2021	AAPC (95% CI)
All ages	Total	21.87	9.41	-6.7 (-8.2, -5.2)*	28.52	11.24	-7.4 (-9.0, -5.7)*	14.34	7.30	-5.6 (-6.9, -4.2)*
	Urban area	14.50	8.21	-4.6 (-6.5, -2.6)*	18.62	9.61	-5.3 (-7.4, -3.1)*	9.97	6.61	-3.4 (-5.1, -1.7)*
	Rural area	27.59	11.79	-6.9 (-8.4, -5.5)*	36.14	14.53	-7.4 (-9.1, -5.8)*	17.75	8.56	-6.0 (-7.4, -4.7)*
Under 1 year	Total	34.37	16.89	-7.9 (-9.7, -6.1)*	41.05	16.82	-8.0 (-9.4, -6.6)*	27.15	16.96	-7.7 (-10.5, -4.8)*
	Urban area	32.63	15.04	-7.2 (-9.7, -4.6)*	36.21	15.19	-7.0 (-9.7, -4.2)*	28.74	14.87	-7.5 (-10.5, -4.5)*
	Rural area	35.50	20.07	-5.3 (-9.0, -1.4)*	44.15	19.62	-7.5 (-8.8, -6.2)*	26.10	20.58	-3.5 (-13.0, 7.0)
1–4 years	Total	53.18	9.16	-11.6 (-14.1, -9.0)*	66.00	10.41	-11.9 (-14.7, -9.0)*	38.40	7.79	-11.0 (-13.1, -8.8)*
	Urban area	27.96	7.68	-8.8 (-11.1, -6.4)*	33.42	8.51	-9.0 (-11.5, -6.4)*	21.93	6.79	-8.4 (-10.4, -6.2)*
	Rural area	71.29	12.18	-11.7 (-14.6, -8.6)*	89.24	14.28	-11.9 (-15.2, -8.5)*	50.31	9.82	-12.8 (-15.9, -9.7)*
5–9 years	Total	12.24	5.44	-7.5 (-9.6, -5.3)*	16.45	6.65	-8.3 (-10.3, -6.3)*	7.53	4.05	-5.9 (-8.7, -2.9)*
	Urban area	8.78	4.68	-3.9 (-8.6, 1.0)	11.29	5.80	-4.5 (-9.3, 0.6)	6.00	3.41	-4.3 (-7.6, -0.8)*
	Rural area	14.90	6.78	-8.0 (-10.4, -5.6)*	20.42	8.14	-9.0 (-11.0, -6.9)*	8.71	5.16	-6.1 (-9.2, -2.8)*
10–14 years	Total	8.80	9.10	0.5 (-2.1, 3.2)	12.49	11.08	-0.8 (-3.6, 2.1)	4.67	6.75	3.0 (-2.0, 8.2)
	Urban area	7.08	7.62	0.7 (-2.8, 4.4)	10.02	8.99	-0.6 (-3.5, 2.3)	3.85	6.03	2.3 (-1.3, 6.2)
	Rural area	10.15	11.32	1.1 (-1.7, 3.9)	14.45	14.21	-0.8 (-4.9, 3.5)	5.32	7.83	3.3 (-1.2, 8.0)*
15–19 years	Total	18.72	13.73	-3.6 (-5.9, -1.3)*	26.26	17.03	-4.8 (-7.1, -2.5)*	10.22	9.96	-0.9 (-3.8, 2.1)
	Urban area	15.14	12.67	-2.1 (-4.8, 0.7)	21.54	15.19	-3.5 (-6.0, -1.0)*	8.17	9.79	1.1 (-2.6, 5.0)
	Rural area	22.90	16.91	-2.2 (-8.5, -4.6)	31.67	22.28	-3.6 (-7.2, 0.2)	12.66	10.52	-2.1 (-4.0, -0.1)*

Abbreviation: APC=annual percent change; AAPC=average annual percent change; CI=confidence interval.

* $P < 0.05$.

SUPPLEMENTARY TABLE S4. APC and AAPC in cause-specific injury mortality among Chinese children and adolescents from 2010 to 2021.

Injury cause	Age group	Area	Block 1		Block 2		Block 3		AAPC (95% CI)
			Time period	APC (95% CI)	Time period	APC (95% CI)	Time period	APC (95% CI)	
Road traffic injury	0–19 years	Urban area	2010–2014	17.3 (–1.5, 39.6)	2014–2021	–10.6 (–16.9, –3.7)*			–1.3 (–7.5, –5.4)
		Rural area	2010–2014	9.1 (–6.3, 27.0)	2014–2021	–10.2 (–15.8, –4.3)*			–3.6 (–9.0, 2.0)
	Under 1 year	Urban area	2010–2021	–8.9 (–18.0, 1.3)					–8.9 (–18.0, 1.3)
		Rural area	2010–2021	–5.6 (–13.3, 2.7)					–5.6 (–13.3, 2.7)
	1–4 years	Urban area	2010–2021	–4.3 (–8.9, 0.7)					–4.3 (–8.9, 0.7)
		Rural area	2010–2021	–7.7 (–10.8, –4.5)*					–7.7 (–10.8, –4.5)*
	5–9 years	Urban area	2010–2014	19.0 (–2.0, 44.5)	2014–2021	–9.7 (–16.8, –1.9)*			–0.1 (–7.2, 7.4)
		Rural area	2010–2012	32.2 (–1.0, 76.5)	2012–2021	–7.5 (–9.9, –5.0)*			–1.3 (–5.8, 3.5)
	10–14 years	Urban area	2010–2014	35.4 (10.8, 65.5)*	2014–2021	–11.3 (–18.5, –3.5)*			3.4 (–4.1, 11.5)
		Rural area	2010–2014	36.0 (16.2, 59.1)*	2014–2021	–9.3 (–15.1, –3.1)*			5.1 (–0.9, 11.5)
	15–19 years	Urban area	2010–2013	21.3 (–1.0, 48.7)	2013–2021	–9.0 (–12.9, –4.9)*			–1.6 (–6.7, 3.8)
		Rural area	2010–2014	13.7 (–6.3, 38.0)	2014–2021	–12.4 (–19.3, –5.0)*			–3.7 (–1.04, 3.6)
Drowning	0–19 years	Urban area	2010–2021	–8.0 (–10.1, –5.8)*					–8.0 (–10.1, –5.8)*
		Rural area	2010–2021	–9.5 (–11.4, –7.5)*					–9.5 (–11.4, –7.5)*
	Under 1 year	Urban area	2010–2017	3.6 (–11.8, 21.7)	2017–2021	–31.1 (–52.9, 0.9)*			–10.7 (–22.6, 3.1)
		Rural area	2010–2021	–13.5 (–17.2, –9.6)*					–13.5 (–17.2, –9.6)*
	1–4 years	Urban area	2010–2021	–12.9 (–15.4, –10.0)*					–12.9 (–15.4, –10.0)*
		Rural area	2010–2012	–37.4 (–45.3, –28.3)*	2012–2021	–11.4 (–12.5, –10.3)*			–16.8 (–18.6, –15.0)*
	5–9 years	Urban area	2010–2013	9.6 (–9.2, 32.3)	2013–2021	–12.8 (–16.3, –9.2)*			–7.2 (–11.7, –2.5)*
		Rural area	2010–2012	32.2 (–1.0, 76.5)	2012–2021	–7.5 (–9.9, –5.0)*			–10.8 (–13.3, –8.1)*
	10–14 years	Urban area	2010–2012	32.5 (4.0, 68.8)*	2012–2021	–9.1 (–11.1, –7.1)*			–2.7 (–6.4, 1.2)
		Rural area	2010–2014	11.0 (–2.4, 26.2)	2014–2019	–12.2 (–22.8, –0.2)*	2019–2021	5.7 (–29.5, 58.6)	–1.1 (–8.2, 6.5)
	15–19 years	Urban area	2010–2013	15.0 (3.0, 28.5)*	2013–2017	–9.2 (–18.7, 1.4)	2017–2021	6.6 (–0.6, 14.3)	2.7 (–1.3, 6.8)
		Rural area	2010–2013	16.9 (–4.6, 43.2)	2013–2018	–9.9 (–20.8, 2.4)	2018–2021	10.6 (–9.7, 35.4)	2.3 (–4.5, 9.6)


Continued

Injury cause	Age group	Area	Block 1		Block 2		Block 3	
			Time period	APC (95% CI)	Time period	APC (95% CI)	Time period	APC (95% CI)
Suffocation	0–19 years	Urban area	2010–2021	-6.2 (-8.5, -3.8)*				-6.2 (-8.5, -3.8)*
		Rural area	2010–2012	18.4 (-7.1, 50.9)	2012–2021	-9.6 (-11.6, -7.6)*		-5.1 (-8.8, -1.3)*
	Under 1 year	Urban area	2010–2021	-7.7 (-10.8, -4.5)*				-7.7 (-10.8, -4.5)*
		Rural area	2010–2012	14.1 (-15.1, 53.2)	2012–2021	-10.9 (-13.3, -8.5)*		-6.8 (-11.2, -2.2)*
	1–4 years	Urban area	2010–2021	-8.7 (-13.1, -4.0)*				-8.7 (-13.1, -4.0)*
		Rural area	2010–2021	-6.0 (-9.3, -2.6)*				-6.0 (-9.3, -2.6)*
	5–9 years	Urban area	2010–2017	17.3 (-3.5, 42.6)	2017–2021	-31.9 (-57.1, 8.1)		-3.7 (-19.1, 14.5)
		Rural area	2010–2021	-2.0 (-8.7, 5.2)				-2.0 (-8.7, 5.2)
	10–14 years	Urban area	2010–2021	4.1 (-6.0, 15.1)				4.1 (-6.0, 15.1)
		Rural area	2010–2021	-0.6 (-10.5, 10.5)				-0.6 (-10.5, 10.5)
Suicide	15–19 years	Urban area	2010–2021	0.7 (-8.0, 10.3)				0.7 (-8.0, 10.3)
		Rural area	2010–2021	NA				NA
	0–19 years	Urban area	2010–2017	1.7 (-2.3, 5.8)	2017–2021	19.9 (9.2, 31.8)*		8.0 (4.2, 11.8)*
		Rural area	2010–2013	13.6 (-5.1, 36.1)	2013–2018	-3.1 (-13.5, 8.7)	2018–2021	18.9 (-0.7, 42.3)
	10–14 years	Urban area	2010–2021	11.5 (7.5, 15.7)*				11.5 (7.5, 15.7)*
		Rural area	2010–2021	11.0 (4.8, 17.6)*				11.0 (4.8, 17.6)*
	15–19 years	Urban area	2010–2017	-2.3 (-9.3, 5.3)	2017–2021	25.8 (5.5, 50.0)*		7.1 (0.3, 14.4)*
		Rural area	2010–2013	7.8 (0.0, 16.2)*	2013–2018	-3.2 (-7.7, 1.5)	2018–2021	15.8 (7.4, 24.8)*
								4.7 (2.1, 7.4)*

Note: NA: Because the initial value is 0, we cannot calculate the APC and AAPC.

Abbreviation: APC=annual percent change; AAPC=average annual percent change; CI=confidence interval; NA=not applicable.

* $P < 0.05$.



SUPPLEMENTARY FIGURE S1. Distribution of injury-related mortality by area and age group in Chinese children and adolescents, from 2010 to 2021. (A) Injury-related mortality distribution in children and adolescents aged 0–19 in urban areas. (B) Injury-related mortality distribution in children and adolescents aged 0–19 in rural areas. (C) Injury-related mortality distribution in children under 1 year in urban areas. (D) Injury-related mortality distribution in children under 1 year in rural areas. (E) Injury-related mortality distribution in children aged 1–4 in urban areas. (F) Injury-related mortality distribution in children aged 1–4 in rural areas. (G) Injury-related mortality distribution in children aged 5–9 in urban areas. (H) Injury-related mortality distribution in children aged 5–9 in rural areas. (I) Injury-related mortality distribution in adolescents aged 10–14 in urban areas. (J) Injury-related mortality distribution in adolescents aged 10–14 in rural areas. (K) Injury-related mortality distribution in adolescents aged 15–19 in urban areas. (L) Injury-related mortality distribution in adolescents aged 15–19 in rural areas.