

## Vital Surveillances

# Epidemiological Characterization of Dengue Fever — Yunnan Province, China, 2010–2021

Xiaoxiang Zhu<sup>1</sup>; Songwang Wang<sup>1</sup>; Yanfei Li<sup>1</sup>; Yu Cao<sup>1</sup>; Xuemei Su<sup>1,†</sup>; Xiaotao Zhao<sup>2,†</sup>

## ABSTRACT

**Objective:** The goal of this study is to analyze the epidemiological patterns of dengue fever across different districts and counties in Yunnan Province from 2010 to 2021.

**Methods:** In this study, we employed joinpoint regression analysis, spatial autocorrelation analysis, and space-time scan analysis to illustrate the spatio-temporal propagation and demographic influence of dengue fever, using both graphical and tabular presentations to clearly demonstrate the findings.

**Results:** Yunnan Province reported 14,098 cases of dengue fever during the period from 2010 to 2021. Of these, 11,513 cases were caused by local transmission, 2,566 were imported internationally, and 19 were inter-provincial imports. Seasonal trends emerged, revealing a surge in incidences during the summer and autumn months. The sex ratio of male to female cases was 1:0.88, with a significant majority of 82.00% of cases involving individuals belonging to the age group of 15–60. Commercial service workers constituted the most impacted occupational group, forming 20.96% of total cases. A spatio-temporal scan identified significant clustering of dengue fever cases across space and time, with the most pronounced cluster observed in southern Yunnan, primarily between 2015 and 2019.

**Conclusions:** Dengue fever in Yunnan Province manifests as biennial outbreaks, underscoring the necessity for increased surveillance, particularly in counties bordering other regions.

Dengue fever, a mosquito-borne illness, can be caused by any of the four dengue virus serotypes (1). Currently, over 100 tropical and subtropical countries are disease-endemic, thus exposing more than a third of the global population to the risk of infection (2). Since 1978, there have been significant outbreaks in

Chinese Mainland, resulting in 655,324 confirmed cases and 610 deaths until 2008. Initially, clustering was observed in southern coastal provincial-level administrative divisions (PLADs) like Guangdong, Hainan, and Guangxi. Nevertheless, the disease distribution started extending northward and westward to include PLADs such as Fujian, Zhejiang, and Yunnan in the 1990s (3–4). Yunnan encountered its first imported case in 1975 but, notably, no other cases were reported for the ensuing 25 years. Despite the sporadic importations of the disease between 2000 and 2007, a significant shift occurred in 2008 when twelve indigenous cases emerged in five border areas of Zhenkang, Mangshi, Yingjiang, Ruili, and Longchuan, indicating the first occurrence of local transmission in the PLAD. An increasing trend in local dengue outbreaks has been noticed recently, both in terms of frequency and geographical distribution since 2010 (5–6). Thus, this paper seeks to elucidate the changing epidemiological characteristics of dengue fever in Yunnan from 2010 to 2021 by examining three distinct periods of the disease's outbreaks within the PLAD.

## METHODS

This research utilized dengue case data from the China Disease Control and Prevention Information System (CDCIS) covering the period from 2010 to 2021. This encompassed both laboratory-confirmed and clinically-diagnosed cases, based on the Diagnostic Criteria for dengue fever (WS 216-2008). The data captured detailed information including patient demographics, inception date of symptoms, diagnostic method, and for cases contracted internationally, a travel history to dengue-endemic regions within a fortnight preceding the onset of symptoms.

The data were analyzed using R statistical software (version 4.3.1, R Foundation for Statistical Computing, Auckland, New Zealand) for case information organization. Microsoft Excel 2016

(Microsoft, Redmond, USA) was utilized for demographic analysis. Joinpoint statistical software (version 5.0.2, Applications Branch, National Cancer Institute, Bethesda, USA) was used to investigate trends in dengue incidence rates and stage characteristics over the study period. The spatio-temporal spread of dengue fever was visually represented using ArcMap (version 10.7, Environmental Systems Research Institute, Redlands, USA) and SaTScan (version 10.1.3, Information Management Services, Maryland, USA). The statistical test significance level was set at 0.05.

## RESULTS

### Epidemic Overview

From 2010 to 2021, Yunnan Province reported 14,098 cases of dengue infection. Of these, 11,513 were locally transmitted, 2,566 were imported from abroad, and 19 were introduced from other PLADs within China. Upon analyzing the infection trends throughout these years, a pattern of sporadic surges was noted. The years 2013, 2015, 2017, and 2019 saw notably high case counts — 1,538, 1,816, 2,575, and 6,471, respectively. Furthermore, from 2013 to 2019,

the regions of Xishuangbanna, Dehong, and Lincang experienced dengue outbreaks in a biennial pattern.

A demographic analysis of dengue cases in Yunnan Province reveals a male-to-female case ratio of roughly 1:0.88. Case ages span from one month to 94 years old, with the majority of cases (approximately 79.83%) identified in adults aged 15–59 years. Cases in children aged 0–14 years constitute about 7.65% of the total, with seniors aged 60 years and above comprising 12.52%. Commercial service workers (2,955 cases), farmers (2,888 cases), and either unemployed individuals or domestic workers (1,569 cases) were identified as the most affected occupational groups, accounting for 20.96%, 20.49%, and 11.13% of total cases, respectively. Statistical analysis reveals significant yearly variations in the distribution of cases by gender, age, and occupation, as comprehensively detailed in Table 1.

### Temporal Distribution

The annual analysis of dengue case numbers across various districts of Yunnan from 2010 to 2021 has revealed a seasonal pattern, with relatively stable, and fewer cases observed during the first half of each year (2.16%). Throughout this period, imported cases were

TABLE 1. Demographic distribution of dengue fever cases in Yunnan Province, 2010–2021.

Item	2010, n (%)	2011, n (%)	2012, n (%)	2013, n (%)	2014, n (%)	2015, n (%)	2016, n (%)	2017, n (%)	2018, n (%)	2019, n (%)	2020, n (%)	2021, n (%)	Total, n (%)	P
Gender														<0.05*
Male	10 (0.13)	12 (0.16)	18 (0.24)	731 (9.73)	136 (1.81)	927 (12.34)	191 (2.54)	1,298 (17.28)	409 (5.44)	3,618 (48.16)	151 (2.01)	11 (0.15)	7,512 (53.28)	
Female	8 (0.12)	1 (0.02)	6 (0.09)	807 (12.25)	77 (1.17)	889 (13.50)	134 (2.03)	1,277 (19.39)	433 (6.57)	2,853 (43.32)	99 (1.50)	2 (0.03)	6,586 (46.72)	
Age														<0.05*
0–14	0 (0.00)	1 (0.09)	1 (0.09)	114 (10.58)	17 (1.58)	95 (8.81)	32 (2.97)	241 (22.36)	73 (6.77)	482 (44.71)	20 (1.86)	2 (0.19)	1,078 (7.65)	
15–59	17 (0.15)	12 (0.11)	22 (0.20)	1,257 (11.17)	189 (1.68)	1,529 (13.59)	269 (2.39)	2,122 (18.85)	681 (6.05)	4,937 (43.86)	209 (1.86)	11 (0.10)	11,255 (79.83)	
≥60	1 (0.06)	0 (0.00)	1 (0.06)	167 (9.46)	7 (0.40)	192 (10.88)	24 (1.36)	212 (12.01)	88 (4.99)	1,052 (59.60)	21 (1.19)	0 (0.00)	1,765 (12.52)	
Occupation														<0.05*
Commercial service	3 (0.10)	2 (0.07)	7 (0.24)	367 (12.42)	89 (3.01)	429 (14.52)	136 (4.60)	618 (20.91)	197 (6.67)	1,060 (35.87)	44 (1.49)	3 (0.10)	2,955 (20.96)	
Farmer	1 (0.03)	2 (0.07)	4 (0.14)	118 (4.09)	29 (1.00)	347 (12.02)	41 (1.42)	585 (20.26)	134 (4.64)	1,591 (55.09)	34 (1.18)	2 (0.07)	2,888 (20.49)	
Domestic helper	0 (0.00)	0 (0.00)	1 (0.06)	194 (12.36)	13 (0.83)	198 (12.62)	30 (1.91)	294 (18.74)	111 (7.07)	702 (44.74)	25 (1.59)	1 (0.06)	1,569 (11.13)	
Retiree	1 (0.08)	0 (0.00)	0 (0.00)	175 (13.90)	2 (0.16)	135 (10.72)	19 (1.51)	140 (11.12)	96 (7.63)	684 (54.33)	7 (0.56)	0 (0.00)	1,259 (8.93)	
Student	2 (0.19)	2 (0.19)	1 (0.10)	140 (13.58)	17 (1.65)	117 (11.35)	28 (2.72)	212 (20.56)	65 (6.30)	423 (41.03)	23 (2.23)	1 (0.10)	1,031 (7.31)	
Others	11 (0.25)	7 (0.16)	11 (0.25)	544 (12.37)	63 (1.43)	590 (13.42)	71 (1.62)	726 (16.52)	239 (5.44)	2,011 (45.75)	117 (2.66)	6 (0.14)	4,396 (31.18)	
Total	18 (0.13)	13 (0.09)	24 (0.17)	1,538 (10.91)	213 (1.51)	1,816 (12.88)	325 (2.31)	2,575 (18.27)	842 (5.97)	6,471 (45.90)	250 (1.77)	13 (0.09)	14,098	

\*: pass Fisher's exact test.

annually recorded, while there was a notable increase in indigenous cases beginning in 2013. Fundamentally, this twelve-year span can be classified into three phases: the early phase (2010–2012), characterized by sporadic and primarily imported cases; the middle phase (2013–2019), which displayed a consistent growth in imported cases alongside a dramatic surge in indigenous cases, with outbreaks occurring biennially; and the last phase (2020–2021), during which both imported and indigenous cases diminished (Figure 1).

The Joinpoint Regression Program analysis discerned a notable alteration in the trend of dengue fever cases in Yunnan Province in 2019 ( $P<0.05$ ). Over the period 2010 to 2019, there was a substantial surge in dengue cases, yielding an annual percentage change (APC) of 85.25% ( $P<0.05$ ). Contrarily, between 2019 and 2021, a precipitous decline in cases was observed, marked by an APC of -95.12% ( $P<0.05$ ) (Figure 2).

The occurrence of dengue fever in Yunnan Province predominantly coincides with the summer and autumn seasons, particularly the latter months of fall. Conversely, the incidence rates during spring and winter are considerably lower, thereby contributing minimally to the annual case count. Monthly analysis indicates a progressive increase in cases starting from June and July each year, culminating in a peak from August to October, followed by a subsequent decline beginning in November. It's worth noting that the resurgence of imported cases precedes that of indigenous cases, yet the latter surge at a quicker pace and peak earlier, as depicted in Figure 3.

### Spatial Distribution

Between 2010 and 2021, a total of 14,098 cases of

dengue fever were documented in Yunnan Province. Out of the sixteen cities and prefectures in the PLAD, Xishuangbanna, Dehong, and Lincang were most impacted. These locations accounted for 8,640 (61.29%), 3,567 (25.30%), and 1,149 (8.15%) cases, respectively. Considerable cases were also reported in Kunming and Honghe, which accounted for 302 (2.14%) and 136 (0.96%) cases, respectively. Dengue fever was reported annually in Kunming and every year, except in 2010, in Dehong, whilst Xishuangbanna reported cases each year except 2021. Dengue fever cases were less prevalent in Baoshan, Zhaotong, Pu'er, Qujing, Dali, Chuxiong, Yuxi, Wenshan, Lijiang, and Nujiang, as these locations all reported fewer than 100 cases. Wenshan has reported cases intermittently since 2018, Lijiang during provincial-wide outbreak peak years, and Nujiang reported a single case in 2017. However, no cases of dengue fever have been reported in Diqing. A detailed account of cases can be seen in Supplementary Table S1 (available at <https://weekly.chinacdc.cn/>).

Between 2010 and 2021, dengue fever was identified in 98 out of 129 districts and counties in Yunnan Province. Of these, 11,513 were indigenous cases spread across 10 districts and counties. Additionally, the PLAD reported 2,566 cases of dengue fever imported from other countries, impacting 96 counties and cities across 15 prefectures. The imported cases primarily originated from Southeast Asian and African countries.

The districts with a significantly high number of indigenous cases were Jinghong (7,571 cases), Ruili (2,620 cases), and Mengla (650 cases), which are all border counties. In terms of imported cases, the top




FIGURE 1. Annual distribution of dengue fever cases in Yunnan Province, 2010–2021.




FIGURE 2. Progression of total dengue fever incidence in Yunnan Province, 2010–2021.

Abbreviation: APC=Annual Percent Change.

\* Indicates that the Annual Percent Change (APC) is significantly different from zero at the  $\alpha = 0.05$  level.

Note: Test statistic and *P*-value not available for the empirical quantile method. Final selected model: 1 Joinpoint.

three districts were Ruili (641 cases), Zhenkang (312 cases), and Gengma (299 cases) as outlined in [Supplementary Table S2](https://weekly.chinacdc.cn/) (available at <https://weekly.chinacdc.cn/>).

Most of the imported cases were from Southeast Asian countries, accounting for 96.96% of all imported cases. Myanmar reported the highest number of cases (1,937 cases), followed by Cambodia (267 cases), Laos (199 cases), Thailand (62 cases), and Vietnam (23 cases). Both Myanmar and Laos have been regular sources of imported cases since 2010, while Cambodia started reporting cases from 2016 onward.

It is noteworthy to mention that Myanmar reported an increase in the number of cases in 2015, 2017, and 2019. Also, 2019 witnessed an increase in reported cases from Laos and Cambodia as detailed in [Supplementary Table S3](https://weekly.chinacdc.cn/) (available at <https://weekly.chinacdc.cn/>).

Furthermore, Yunnan Province recorded 19 cases of dengue fever imported from other PLADs within China, predominantly Guangdong. The majority of these cases were reported in Jinghong, Guandu, and Ruili, with particular spikes in the years 2014 and 2019.

### Spatio-Temporal Aggregation

**Global spatial autocorrelation analysis:** The analysis of global spatial autocorrelation for dengue incidence in Yunnan PLAD suggested significant spatial

clustering in specific years. Moran's *I* values were 0.427 for both 2010 and 2012, followed by 0.05, 0.016, 0.172, 0.107, and 0.128 for 2013, 2016, 2017, 2018, and 2019 correspondingly. These values, with statistical significance ( $P < 0.05$ ), indicated considerable spatial clustering of dengue cases. Nevertheless, Moran's *I* value for 2014 was -0.002 ( $P < 0.05$ ), implying a random distribution of cases for that year. During other years, represented by Moran's *I* values of 0.233, 0.008, and -0.003, the differences were not statistically significant ( $P > 0.05$ ), suggesting no noteworthy spatial clustering. Further details can be found in [Supplementary Table S4](https://weekly.chinacdc.cn/) (available at <https://weekly.chinacdc.cn/>).

**Local spatial autocorrelation analysis:** From 2010 to 2021, each year saw the identification of 1 to 5 dengue fever hotspots within the counties of Yunnan PLAD. In the span of 2010 to 2012, these hotspots, largely comprising imported cases, were primarily found in locations such as Jinghong, Ruili, and Kunming. However, between 2013 and 2021, the principal concentration of dengue fever cases transitioned to predominantly include Jinghong and Ruili, with the spread of the disease radiating outward from these two central hotspots. These findings are detailed in [Supplementary Table S5](https://weekly.chinacdc.cn/) (available at <https://weekly.chinacdc.cn/>).

**Spatio-temporal scan analysis:** The spatiotemporal cluster analysis of dengue fever in Yunnan uncovered




FIGURE 3. Monthly trends in dengue fever cases in Yunnan Province, 2010–2021.

distinctive temporal and spatial patterns. This analysis pinpointed a primary cluster and two secondary clusters via spatial and temporal scan techniques (Supplementary Table S6, available at <https://weekly.chinacdc.cn/>). The principle cluster was located in the southern quadrant of the PLAD. Its highest-risk phase spanned from 2015 to 2019, marked by a relative risk (RR) of 150.46, a log likelihood ratio (LLR) of 25770.59, and a highly significant  $P$ -value ( $P < 0.001$ ). This high-risk cluster, which covered a circular area with a radius of 69.29 km and included the counties of Jinghong and Mengla, reported a total of 7,085 dengue fever cases.

## CONCLUSIONS

Dengue fever in Yunnan Province is primarily influenced by imported cases, predominantly from Myanmar and Laos, which often result in local outbreaks. This assertion is bolstered by various studies that underscore imported cases as the principal cause of local epidemics, aligning with the results of our research (7–9). It is advisable for authorities to diligently observe dengue fever trends in Southeast Asian countries, specifically Myanmar and Laos, during the year's first half. It is imperative to intensify screening procedures at Yunnan's borders, particularly

at points of entry.

The border regions of Yunnan have been identified as significant hotspots for transmission, primarily attributed to the *Aedes aegypti* mosquito. Over the last decade, the number of affected districts and counties has surged from nine in 2010 to 98 in 2021. This substantial increase may be associated with the widespread presence of the *Aedes aegypti* mosquito and the mobility of infected individuals (11–12). The high density of mosquito populations along the borders, coupled with diverse vector species, coincides with the seasonal spikes of dengue in the PLAD (13–14). It is recommended to continue surveillance of the *Aedes aegypti* mosquito populations, ensure efficient mosquito control, and uphold cleanliness in border areas. Continuous, year-round monitoring is crucial in key regions such as Ruili, Jinghong, and Gengma. Additionally, public health initiatives should drive sanitation practices to prevent the creation of mosquito breeding sites.

There was a discernible alteration in the patterns of dengue occurrence in Yunnan after 2019. This change may be attributed to various factors including the implementation of COVID-19 control measures, enhanced cooperative preventive initiatives, more stringent entry and isolation policies for foreign arrivals, amplified screening and regulation of



movement, the endorsement of remote work, the practice of social distancing, and a decline in public mobility. Collectively, these factors led to a substantial decrease in reported dengue cases.

In conclusion, Yunnan Province has seen sporadic outbreaks of dengue fever, generally on a biennial cycle. Vigilance remains essential regarding the potential increase in prevalence of dengue fever in border counties. Implementing effective dengue surveillance and quarantine measures for travelers originating from regions endemic with the disease is of utmost importance, particularly during large outbreaks or seasons of elevated dengue risk.

Dengue fever in Yunnan Province demonstrates biennial outbreaks, which emphasizes the importance of increased surveillance of the disease in counties located near the border.

**Funding:** National Science and Technology Infrastructure Platform, National Population and Health Science Data Sharing Service Platform, Public Health Science Data Center (NCMI-ZB01N-201905).

doi: 10.46234/ccdcw2024.088

# Corresponding authors: Xuemei Su, [suxm@chinacdc.cn](mailto:suxm@chinacdc.cn); Xiaotao Zhao, [zxt\\_423@163.com](mailto:zxt_423@163.com).

<sup>1</sup> National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, Chinese Center for Disease Control and Prevention, Beijing, China; <sup>2</sup> Yunnan Institute of Parasitic Diseases, Yunnan, China.

Submitted: January 15, 2024; Accepted: May 09, 2024

## REFERENCES

- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and burden of dengue. *Nature* 2013;496(7446):504–7. <https://doi.org/10.1038/nature12060>.
- Brady OJ, Gething PW, Bhatt S, Messina JP, Brownstein JS, Hoen AG, et al. Refining the global spatial limits of dengue virus transmission by evidence-based consensus. *PLoS Negl Trop Dis* 2012;6(8):e1760. <https://doi.org/10.1371/journal.pntd.0001760>.
- Wu JY, Lun ZR, James AA, Chen XG. Dengue Fever in mainland China. *Am J Trop Med Hyg* 2010;83(3):664–71. <https://doi.org/10.4269/ajtmh.2010.09-0755>.
- Yue YJ, Liu XB, Xu M, Ren DS, Liu QY. Epidemiological dynamics of dengue fever in mainland China, 2014–2018. *Int J Infect Dis* 2019;86:82–93. <https://doi.org/10.1016/j.ijid.2019.06.015>.
- Yue YJ, Liu QY, Liu XB, Zhao N, Yin WW. Dengue fever in mainland China, 2005–2020: a descriptive analysis of dengue cases and *Aedes* data. *Int J Environ Res Public Health* 2022;19(7):3910. <https://doi.org/10.3390/ijerph19073910>.
- Li HX, Zhou HN, Yang YC, Jiang H. Dengue fever epidemic situation in Yunnan province from 2004 to 2008. *Chin J Vector Biol Control* 2010;21(6):576–7,580. <https://d.wanfangdata.com.cn/periodical/zgmjswxjkzzz201006017>. (In Chinese).
- Wang BH, Yang HL, Feng Y, Zhou HN, Dai JJ, Hu YZ, et al. The distinct distribution and phylogenetic characteristics of dengue virus serotypes/genotypes during the 2013 outbreak in Yunnan, China: Phylogenetic characteristics of 2013 dengue outbreak in Yunnan, China. *Infect Genet Evol* 2016;37:1–7. <https://doi.org/10.1016/j.meegid.2015.10.022>.
- Wang BH, Li YP, Feng Y, Zhou HN, Liang YB, Dai JJ, et al. Phylogenetic analysis of dengue virus reveals the high relatedness between imported and local strains during the 2013 dengue outbreak in Yunnan, China: a retrospective analysis. *BMC Infect Dis* 2015;15:142. <https://doi.org/10.1186/s12879-015-0908-x>.
- Hu TS, Zhang HL, Feng Y, Fan JH, Tang T, Liu YH, et al. Epidemiological and molecular characteristics of emergent dengue virus in Yunnan Province near the China-Myanmar-Laos border, 2013–2015. *BMC Infect Dis* 2017;17(1):331. <https://doi.org/10.1186/s12879-017-2401-1>.
- Zhang J, Shu Y, Shan XY, Li DY, Ma DH, Li TT, et al. Co-circulation of three dengue virus serotypes led to a severe dengue outbreak in Xishuangbanna, a border area of China, Myanmar, and Laos, in 2019. *Int J Infect Dis* 2021;107:15–7. <https://doi.org/10.1016/j.ijid.2021.04.010>.
- Liu YH, Yin XX, Zhang HL, Yang ZL, Dong CL, Chen SY, et al. Epidemiological characteristics of dengue fever and monitoring of *Aedes* vector mosquitoes in Dehong Dai and Jingpo Autonomous Prefecture of Yunnan province, China, 2013–2019. *Chin J Vector Biol Control* 2021;32(2):173–80. <https://doi.org/10.11853/j.issn.1003.8280.2021.02.011>.
- Zhang HL, Zhang YZ, Yang WH, Feng Y, Nasci RS, Yang J, et al. Mosquitoes of Western Yunnan Province, China: seasonal abundance, diversity, and arbovirus associations. *PLoS One* 2013;8(10):e77017. <https://doi.org/10.1371/journal.pone.0077017>.
- Liang GD, Li XL, Gao XY, Fu SH, Wang HY, Li MH, et al. Arboviruses and their related infections in China: a comprehensive field and laboratory investigation over the last 3 decades. *Rev Med Virol* 2018;28(1):e1959. <https://doi.org/10.1002/rmv.1959>.
- Lv RC, Zhu CQ, Wang CH, Ai LL, Lv H, Zhang B, et al. Genetic diversity and population structure of *Aedes aegypti* after massive vector control for dengue fever prevention in Yunnan border areas. *Sci Rep* 2020;10(1):12731. <https://doi.org/10.1038/s41598-020-69668-7>.
- Li NZ, Feng Y, Vrancken B, Chen YY, Dong L, Yang QQ, et al. Assessing the impact of COVID-19 border restrictions on dengue transmission in Yunnan Province, China: an observational epidemiological and phylogenetic analysis. *Lancet Reg Health West Pac* 2021;14:100259. <https://doi.org/10.1016/j.lanwpc.2021.100259>.

## SUPPLEMENTARY MATERIAL

SUPPLEMENTARY TABLE S1. Distribution of dengue fever cases across prefectures and cities in Yunnan Province, China, 2010–2021.

City	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	Total	Component ratio (%)
Xishuangbanna	6	4	14	1327	1	1141	32	1356	637	4126	1	0	8645	61.32
Dehong	0	1	1	193	188	448	279	837	98	1271	240	11	3567	25.30
Lincang	0	0	0	0	1	196	1	319	43	587	2	0	1149	8.15
Kunming	8	7	5	14	15	13	11	22	34	166	4	2	301	2.14
Honghe	0	0	0	0	0	3	0	5	2	125	1	0	136	0.96
Baoshan	0	0	0	2	3	3	1	23	4	38	0	0	74	0.52
Zhaotong	0	0	0	0	2	1	0	4	10	43	1	0	61	0.43
Puer	3	0	2	0	0	3	1	2	2	45	0	0	58	0.41
Qujing	1	1	1	0	1	2	0	3	3	29	0	0	41	0.29
Dali	0	0	0	1	1	2	0	1	4	19	0	0	28	0.20
Chuxiong	0	0	0	0	1	0	0	2	2	8	0	0	13	0.09
Yuxi	0	0	0	0	0	3	0	0	2	6	0	0	11	0.08
Wenshan	0	0	0	0	0	0	0	0	1	5	1	0	7	0.05
Lijiang	0	0	0	1	0	1	0	0	0	3	0	0	5	0.04
Nujiang	0	0	0	0	0	0	0	1	0	0	0	0	1	0.01
Diqing	0	0	0	0	0	0	0	0	0	0	0	0	0	0.00
Unknown	0	0	1	0	0	0	0	0	0	0	0	0	1	0.01
Total	18	13	24	1538	213	1816	325	2575	842	6471	250	13	14098	100.00

SUPPLEMENTARY TABLE S2. Dengue fever cases by district and county in Yunnan Province, 2010–2021.

County	Cumulative cases	Indigenous cases	Overseas imported cases	Imported cases from other PLADs
Wuhua District	48	0	48	0
Panlong District	43	0	43	0
Guandu District	115	0	112	3
Xishan District	49	0	48	1
Dongchuan District	4	0	4	0
Chenggong District	9	0	9	0
Jinning District	2	0	2	0
Fumin County	3	0	3	0
Yiliang County	6	0	6	0
Shilin Yi Autonomous County	1	0	1	0
Songming County	7	0	7	0
Luchuan Yi and Miao Autonomous County	4	0	4	0
Xundian Hui and Yi Autonomous County	2	0	2	0
Anning City	4	0	4	0
Qilin District	6	0	6	0
Zhanyi District	2	0	2	0
Malong District	0	0	0	0
Luliang County	3	0	3	0
Shizong County	3	0	3	0

Continued

County	Cumulative cases	Indigenous cases	Overseas imported cases	Imported cases from other PLADs
Luoping County	5	0	5	0
Fuyuan County	4	0	3	1
Huize County	6	0	6	0
Xuanwei City	12	0	12	0
Hongta District	2	0	2	0
Jiangchuan District	3	0	3	0
Tonghai County	2	0	2	0
Huaning County	0	0	0	0
Yimen County	1	0	1	0
Eshan Yi Autonomous County	1	0	1	0
Xinping Yi and Dai Autonomous County	0	0	0	0
Yuanjiang Hani, Yi, and Dai Autonomous County	0	0	0	0
Chengjiang City	2	0	2	0
Longyang District	12	0	12	0
Shidian County	6	0	6	0
Longling County	11	0	11	0
Changning County	0	0	0	0
Tengchong City	45	0	44	1
Zhaoyang District	4	0	4	0
Ludian County	3	0	3	0
Qiaojia County	2	0	2	0
Yanjin County	4	0	4	0
Daguan County	0	0	0	0
Yongshan County	2	0	2	0
Suijiang County	1	0	1	0
Zhenxiong County	41	0	40	1
Yiliang County	1	0	1	0
Weixin County	2	0	2	0
Shuifu City	1	0	1	0
Gucheng District	0	0	0	0
Yulong Naxi Autonomous County	2	0	2	0
Yongsheng County	3	0	2	1
Huaping County	0	0	0	0
Ninglang Yi Autonomous County	0	0	0	0
Simao District	8	0	8	0
Ninger Hani and Yi Autonomous County	0	0	0	0
Mojiang Hani Autonomous County	0	0	0	0
Jingdong Yi Autonomous County	3	0	3	0
Jinggu Dai and Yi Autonomous County	2	0	2	0
Zhenyuan Yi, Hani and Lahu Autonomous County	0	0	0	0
Jiangcheng Hani and Yi Autonomous County	5	3	2	0
Menglian Dai, Lahu and Wa Autonomous County	37	15	22	0



Continued

County	Cumulative cases	Indigenous cases	Overseas imported cases	Imported cases from other PLADs
Lancang Lahu Autonomous County	2	0	2	0
Ximeng Wa Autonomous County	1	0	1	0
Linxiang District	4	0	4	0
Fengqing County	7	0	7	0
Yun County	3	0	3	0
Yongde County	12	0	12	0
Zhenkang County	312	0	312	0
Shuangjiang Lahu, Wa, Bulang, and Dai Autonomous County	2	0	2	0
Gengma Dai and Wa Autonomous County	802	502	299	1
Cangyuan Wa Autonomous County	7	0	7	0
Chuxiong City	3	0	3	0
Shuangbai County	0	0	0	0
Mouding County	4	0	4	0
Nanhua County	1	0	0	1
Yao'an County	2	0	2	0
Dayao County	1	0	1	0
Yongren County	0	0	0	0
Yuanmou County	0	0	0	0
Wuding County	1	0	1	0
Lufeng County	1	0	1	0
Gejiu City	1	0	1	0
Kaiyuan City	3	0	3	0
Mengzi City	3	0	3	0
Mile City	5	0	5	0
Pingbian Miao Autonomous County	1	0	1	0
Jianshui County	0	0	0	0
Shiping County	0	0	0	0
Luxi County	1	0	1	0
Yuanyang County	1	0	1	0
Honghe County	0	0	0	0
Jinping Miao, Yao and Dai Autonomous County	3	0	3	0
Luchun County	1	0	1	0
Hekou Yao Autonomous County	117	114	3	0
Wenshan City	1	0	1	0
Yanshan County	3	0	3	0
Xichou County	0	0	0	0
Malipo County	0	0	0	0
Maguan County	0	0	0	0
Qiubei County	0	0	0	0
Guangnan County	2	0	1	1
Funing County	1	0	0	1
Jinghong City	7,668	7,571	95	2

Continued

County	Cumulative cases	Indigenous cases	Overseas imported cases	Imported cases from other PLADs
Menghai County	211	0	211	0
Mengla County	765	650	115	0
Dali City	5	0	5	0
Yangbi Yi Autonomous County	0	0	0	0
Xiangyun County	6	0	6	0
Binchuan County	3	0	2	1
Midu County	2	0	2	0
Nanjian Yi Autonomous County	2	0	2	0
Weishan Yi and Hui Autonomous County	1	0	1	0
Yongping County	0	0	0	0
Yunlong County	2	0	2	0
Eryuan County	6	0	6	0
Jianchuan County	1	0	1	0
Heqing County	0	0	0	0
Ruili City	3,262	2,620	641	3
Mang City	51	18	33	0
Lianghe County	10	0	7	1
Yingjiang County	161	7	154	0
Longchuan County	83	13	70	0
Lushui City	0	0	0	0
Fugong County	0	0	0	0
Gongshan Derung and Nu Autonomous County	0	0	0	0
Lanping Bai and Pumi Autonomous County	1	0	1	0
Shanggelila City	0	0	0	0
Deeqeen County	0	0	0	0
Weixi Lisu Autonomous County	0	0	0	0
Unknown	6	0	6	0
Total	14,098	11,513	2,566	19

SUPPLEMENTARY TABLE S3. Countries of origin and number of dengue cases imported into Yunnan Province, 2010–2021.

City	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	Total	Component ratio (%)
Myanmar	1	2	2	65	59	285	91	618	101	707	5	1	1,937	75.49
Cambodia	0	0	0	0	0	0	1	0	42	222	2	0	267	10.41
Laos	7	4	13	14	0	5	5	32	2	115	1	1	199	7.76
Thailand	0	0	3	8	2	10	5	4	12	18	0	0	62	2.42
Africa	0	2	1	3	7	1	2	1	0	8	2	0	27	1.05
Vietnam	0	0	0	0	0	2	0	5	2	12	2	0	23	0.90
Malaysia	0	0	0	0	1	2	2	1	0	3	0	0	9	0.35
Sri Lanka	0	0	0	0	0	2	1	2	2	0	0	1	8	0.31
India	1	0	2	0	0	0	0	1	0	2	0	0	6	0.23
Bangladesh	0	0	0	0	0	1	0	1	0	1	0	0	3	0.12
Maldives	0	0	0	0	0	0	0	0	2	1	0	0	3	0.12
Philippines	0	0	0	0	0	0	0	0	0	2	1	0	3	0.12
Indonesia	0	0	0	0	0	0	1	0	0	0	0	0	1	0.04
Unknown	9	5	3	1	0	0	0	0	0	0	0	0	18	0.70
Total	18	13	24	91	69	308	108	665	163	1,091	13	3	2,566	100.00

SUPPLEMENTARY TABLE S4. Spatial global autocorrelation analysis of dengue fever in Yunnan Province, China, 2010–2021.

Year	Moran's I	Z	P
2010	0.427	9.019	<0.001
2011	0.019	0.556	0.527
2012	0.581	13.761	<0.001
2013	0.05	2.010	0.044
2014	-0.002	2.249	0.025
2015	0.008	0.389	0.697
2016	0.016	4.079	<0.001
2017	0.172	4.535	<0.001
2018	0.107	3.510	<0.001
2019	0.128	3.240	0.001
2020	0.000074	3.324	<0.001
2021	-0.003	1.403	0.161

SUPPLEMENTARY TABLE S5. Hot spots of dengue fever annual incidence in Yunnan Province, China.

Year	Hotspot	
	Number	County
2010	5	Jinghong, Mengla, Simao, Chenggong, Panlong
2011	5	Jinghong, Ruili, Wuhua, Anning, Jinning
2012	4	Jinghong, Mengla, Jiangcheng, Ruili
2013	2	Jinghong, Ruili
2014	1	Ruili
2015	3	Jinghong, Ruili, Gengma
2016	1	Ruili
2017	4	Jinghong, Ruili, Mengla, Gengma
2018	4	Jinghong, Mengla, Menghai, Simao
2019	3	Jinghong, Mengla, Ruili
2020	1	Ruili
2021	1	Ruili

SUPPLEMENTARY TABLE S6. Analysis of temporal and spatial scanning results of dengue fever incidence in Yunnan Province, China, 2010–2021.

Type	Number	Radius (km)	Time frame	Cases	Expected cases	Population	RR	LLR	P
Most likely cluster	2	69.29	2015–2019	7,085	94.07	732,398	150.46	25770.59	< 0.001
Secondary likely cluster 1	54	277.33	2010–2015	71	3521.99	22,853,680	0.015	3684.24	< 0.001
Secondary likely cluster 2	29	181.40	2017–2019	3,236	732.51	9,510,581	5.44	2554.75	< 0.001

Abbreviation: RR= relative risk; LLR= log-likelihood ratio.