

## Preplanned Studies

## Persistent Urinary Tract Infection in Association with Community-Acquired NDM-5 *Escherichia coli* Clonal Group Following COVID-19 Infection — Beijing Municipality, China, 2023

Jiazhen Guo<sup>1,8</sup>; Ran Duan<sup>2,8</sup>; Dan Zhang<sup>2</sup>; Peng Zhang<sup>2</sup>; Shuai Qin<sup>2</sup>; Yajuan Fang<sup>1</sup>; Yingna Sun<sup>1</sup>; Lianhe Lu<sup>1</sup>; Huaiqi Jing<sup>2</sup>; Xin Wang<sup>2,9</sup>; Rongmeng Jiang<sup>1,9</sup>; Biao Kan<sup>3,9</sup>

### Summary

#### What is already known about this topic?

The hospital-acquired infections caused by New Delhi metallo-beta-lactamase (NDM)-producing strains are typically attributed to a single clonal lineage.

#### What is added by this report?

In this study, we encountered a unique case of community-acquired NDM-5 *Escherichia coli* urinary tract infection (UTI) following coronavirus disease 2019 (COVID-19). The UTI persisted for a duration of at least 45 days. Genomic analyses revealed the presence of two NDM-5 strains, both sharing an identical chromosomal background but distinct, homologous, and recombined plasmids. This case suggests that a diverse range of resistance genes may be present within the human body, with drug-resistant strains undergoing continuous evolution during infection. The intestinal tract may have been its drug-resistant gene pool.

#### What are the implications for public health practice?

The observations presented in this case indicate that the endogenous acquisition of drug-resistant genes may also be an issue in managing multidrug-resistant organisms (MDRO). It is possible for continuous recombination to occur within carbapenem-resistant Enterobacteriaceae (CRE) during infection. In contrast to exogenously-acquired resistance, greater attention should be placed on the endogenous factors that contribute to the development of CRE within healthcare settings.

Carbapenem-resistant Enterobacteriaceae (CRE) infections, specifically those involving New Delhi metallo-beta-lactamase (NDM) drug-resistant bacteria, continue to pose challenges for healthcare institutions. High-risk coronavirus disease 2019 (COVID-19) patients are susceptible to developing secondary bacterial coinfections, with CRE coinfections

reportedly having significant implications for COVID-19 prognosis. In this report, we present a unique case of a NDM-producing bacterial coinfection following COVID-19 infection.

In late December 2022, a hospital in Beijing admitted an elderly female patient who had been infected with COVID-19 for three weeks. Subsequent laboratory tests revealed a persistent urinary tract infection (UTI) lasting over 45 days, caused by community-acquired NDM-5 *Escherichia coli*. Genomic analyses demonstrated that the infection was not the result of a single strain but rather a group of clonally related, chromosomally indistinguishable *E. coli* strains with varying acquisitions or losses of drug-resistance plasmids. During the course of the infection, a novel drug-resistance plasmid recombined within the patient.

This case highlights the continuous evolution of drug-resistant bacterial strains during infections and suggests that the intestinal tract may serve as a source for the gene pool.

A unique case was identified in a hospital in Beijing involving a patient admitted from December 29, 2022, to January 13, 2023. CRE *E. coli* was isolated from both the patient's sputum and urine on December 31, 2022. Medical records and clinical test results were collected for analysis. The patient was followed up twice to isolate the multidrug-resistant organisms (MDRO): urine and feces samples were collected on January 26, 2023, and additional urine, feces, and sputum samples were obtained on February 17, 2023. A urine routine test was also conducted on February 17, 2023. Concurrently, control samples were collected from the patient's live-in nanny on February 17, 2023.

The antibiotic sensitivity of the strains was evaluated using minimum inhibitory concentrations (MICs) testing using VITEK<sup>®</sup> 2 COMPACT (bioMérieux, Marcy-l'Étoile, France), and *bla*<sub>NDM</sub> genes were detected and sequenced in positive samples (1). All strains underwent molecular typing with MLST (2), and whole-genome sequencing was carried out using

PacBio Sequel (Pacific Biosciences, Menlo Park, USA) and Illumina NovaSeq (Illumina, San Diego, USA) technologies. Antibiotic resistance genes were predicted using ResFinder 4.1 (Technical University of Denmark, Kongens Lyngby, Denmark) (3).

Genomic synteny between two plasmids was visualized using ACT release 13.0.0 (Wellcome Trust Sanger Institute, Cambridge, UK). Primers were designed and amplified for all assembled plasmids identified through whole-genome analysis (Supplementary Table S1, available in <https://weekly.chinacdc.cn/>). A phylogenetic tree based on the core chromosomal genome was constructed using the neighbor-joining method (4–6). Additionally, a NJ tree for plasmids and associated megaBLAST hits was developed based on pairwise alignments.

The patient, a 93-year-old woman with a smoking history of over 75 years, tested positive for COVID-19 antigen three weeks prior. In early and mid-December 2022, she experienced a fever that peaked at 38.7 °C, accompanied by mild chills. At that time, she was severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) antigen-positive. Subsequently, her body temperature returned to normal without the use of antipyretics. She self-medicated with a second-

generation cephalosporin antibiotic for three weeks without medical consultation. Around December 20, 2022, she sustained burns to her neck, face, and auricle due to smoking while inhaling oxygen at home. Approximately two days later, she developed a low-grade fever, exhibited a decreased appetite, and showed signs of mental fatigue.

On December 29, 2022, the patient was admitted to the hospital for persistent discomfort following the COVID-19 infection and neck and facial burns. She did not report any urinary tract discomfort during admission or hospitalization and denied any abnormal discharge. On the day of admission, PCR tests for SARS-CoV-2, influenza A, influenza A (H1N1), seasonal influenza A (H3), and influenza B were all negative, and antibody testing for *Mycoplasma pneumoniae* was also negative. As the SARS-CoV-2 infection had resolved at least two weeks prior, Paxlovid was not indicated. The treatment focused on maintaining oxygenation status, managing water and electrolyte balance, and addressing the skin burns.

Due to elevated blood neutrophil levels on admission (Table 1), a bacterial infection was suspected. Consequently, moxifloxacin was added to the treatment regimen at a dosage of 0.4 g

TABLE 1. Significant clinical test results for the patient following COVID-19 infection — Beijing, China, 2023.

Clinical tests	December 29, 2022	December 31, 2022	January 5, 2023	January 9, 2023	February 17, 2023
<b>Blood</b>					
White blood cell	8.95×10 <sup>9</sup>	9.85×10 <sup>9</sup>	5.6×10 <sup>9</sup>	5.81×10 <sup>9</sup>	–
Neutrophil count	7.03×10 <sup>9</sup>	8.49×10 <sup>9</sup>	3.71×10 <sup>9</sup>	3.85×10 <sup>9</sup>	–
Neutrophil (%)	78.50	86.20	66.10	66.20	–
Lymphocyte count	1.03×10 <sup>9</sup>	0.79×10 <sup>9</sup>	1.2×10 <sup>9</sup>	1.15×10 <sup>9</sup>	–
Lymphocyte (%)	11.50	8.00	21.40	19.80	–
Monocytes count	0.82×10 <sup>9</sup>	0.53×10 <sup>9</sup>	0.58×10 <sup>9</sup>	0.62×10 <sup>9</sup>	–
Monocytes (%)	9.20	5.44	10.30	10.70	–
ESR (mm/h)	65	–	74	–	–
CK (U/L)	102.6	280.8	74.3	–	–
CRP (mg/L)	56.3	94.0	8.8	6.4	–
SAA (mg/L)	311.8	392.2	66.9	23.6	–
PCT (ng/L)	<0.05	<0.05	<0.05	<0.05	–
D-Dimer (mg/L)	3.10	–	1.21	1.07	–
<b>Urine</b>					
Leukocytes in the urine (p/HPF)	–	128.77	104.03	14.48	168.83
Red blood cells in the urine (p/HPF)	–	1.40	3.56	5.19	0.96
Bacteria count in the urine (p/HPF)	–	3177.32	912.81	729.26	1002.55

Note: “–” means data not available.

Abbreviation: ESR=erythrocyte sedimentation rate; CK=creatinine kinase; CRP=C reactive protein; SAA=serum amyloid A; PCT=procalcitonin.

intravenously per day, commencing on December 31, 2022, and ceasing on January 6, 2023. The patient was discharged on January 13, 2023, due to improvement in her condition.

CRE was isolated from the patient's urine and sputum on December 31, 2022, from her urine on January 5, 2023, and from her urine and feces on January 26, 2023, and February 17, 2023. No CRE was isolated from the patient's live-in nanny at any time point. All strains isolated from the patient carried the *bla*<sub>NDM-5</sub> gene (accession numbers OQ357728, OQ357729, OQ442328, OQ442329, OQ442330, OQ851640, and OQ851641). The antimicrobial resistance spectra of NDM-5-producing *E. coli* indicated two types, which differed between the strains isolated from urine and sputum on December 31, 2022, with respect to gentamicin, tobramycin, and trimethoprim/sulfamethoxazole (Table 2). To exclude non-CRE *E. coli* infection, the *bla*<sub>NDM</sub> gene positivity rate among 95 *E. coli* isolates acquired from urine on February 17, 2023, was determined to be 88.42%. The antimicrobial resistance results for gentamicin, tobramycin, and trimethoprim/sulfamethoxazole of these NDM-producing *E. coli* isolates were identical (Table 2). The patient was admitted to a new ward that opened on December 21, 2022. No MDRO had been detected in the ward's environment or among other patients from the ward's opening until four weeks after the patient with NDM-5-producing *E. coli* was discharged (December 21, 2022 to February 10, 2023). The patient had not been hospitalized in the past three years, suggesting that the NDM-5-producing *E. coli* was community-acquired.

A group of NDM-5-producing *E. coli* strains were isolated four times from a patient, and they shared the same chromosomes (CP123592, CP123593). Each of these strains carried four plasmids, three of which were identical: pDJH\_Plas\_NDM-5 (CP123590), pDJH\_Plas3 (CP123589), and pDJH\_Plas4 (CP123594). The distinct plasmids were pDJH\_137urine\_Plas1 (CP123591) and pDJH\_183sputum\_Plas2 (CP123595) (Supplementary Table S2, available in <https://weekly.chinacdc.cn/>). Figure 1 displays the timeline of strain isolation, genomic composition, and homology analysis.

All NDM-5 *E. coli* strain sequence types were identified as ST 410, which is distinct from the common uropathogenic *E. coli* (UPEC) strains. The chromosome size was 4.8 Mb, with a G+C content of 50.67 mol%. A NJ tree, including the five sequenced strains and representative strains from each lineage, was divided into two branches. The *E. coli* chromosome in

this study was part of branch C, which was relatively distant from most UPEC strains. The nearest human strain was enteroinvasive *E. coli* (EIEC) with a chromosome size of 4.8 Mb and G+C content of 50.7 mol%.

Common virulence factors of UPEC such as *fimH*, *afa*, *fyuA*, *hlyA*, *iucABCD*, *kpsF*, and *sfa* were found within the chromosome. Notably, type I fimbriae (*fimH*) plays a crucial role in UPEC's ability to colonize the urethral epithelium and migrate to the bladder. Additionally, the chromosome encodes for the locus of enterocyte effacement (LEE) (7–8).

The pDJH\_Plas\_NDM-5 is a 48Kbp IncX3 plasmid that carries the *bla*<sub>NDM-5</sub> gene. It has a similar backbone to *bla*<sub>NDM-5</sub> plasmids of various sources, which range from 46,155 to 53,056 bp in size, with 84% (42/50) of them being 46,161 bp. None of the top 50 megaBlast hits was the same size as Plas\_NDM-5 of 48,020 bp. The extra sequences in this case encoded IS91 family transposase and a putative protein of unknown function. pDJH\_Plas\_NDM-5 plasmid remained stable after 70 passages in vitro.

The pDJH\_183sputum\_Plas2 plasmid (48.6 Kbp) exhibited 31.87% coverage and 99.78% identity when compared with pDJH\_137urine\_Plas1 (152.2 Kbp), suggesting their homology. The top megaBLAST hit demonstrated 96% coverage and 99.87% identity with pDJH\_183sputum\_Plas2. In contrast, the top hit for pDJH\_137urine\_Plas1 revealed only 67% coverage, implying that it was not a pre-existing plasmid but rather a newly emerged plasmid. The pDJH\_183sputum\_Plas2 plasmid encodes the blaTEM-1B gene, which mediates resistance to ampicillin, cephalothin, piperacillin, and ticarcillin. The pDJH\_137urine\_Plas1 plasmid encodes 11 resistance genes that mediate resistance to 30 types of antibiotics across seven categories: sulfonamides, aminoglycosides, phosphomycin, macrolides, tetracyclines, β-lactams, and acyl alcohol.

The differences in drug resistance between strains DJH\_183sputum and DJH\_137urine (Table 2), specifically concerning compound sulfonamides, gentamicin, and tobramycin, were attributed to the presence of *dfrA12*, *sul2*, and *aac(3)-IV*, which were exclusively found in the pDJH\_137urine\_Plas1 plasmid. Therefore, it was deduced that pDJH\_137urine\_Plas1 originated from recombination with pDJH\_183sputum\_Plas2 and acquired new resistance genes within the patient's body. Initially, on December 31, 2022, the patient's urine and sputum strains carried these two plasmids, respectively. On January 5, 2023, the urine strain (JARVKQ0000 00000) harbored pDJH\_137urine\_Plas1. However,

TABLE 2. Spectrum of antibiotic resistance among *bla*<sub>NDM-5</sub> strains isolated from the patient following COVID-19 infection — Beijing, China, 2023.


Antibiotic	December 31, 2022			January 5, 2023			January 26, 2023			February 17, 2023		
	Urine		Sputum	Urine		Feces	Urine		Feces	Urine		Feces
	MIC (μg/mL)	Interpretation	MIC (μg/mL)	Interpretation	MIC (μg/mL)	Interpretation	MIC (μg/mL)	Interpretation	MIC (μg/mL)	Interpretation	MIC (μg/mL)	Interpretation
Ampicillin	≥32	R	≥32	R	≥32	R	≥32	R	≥32	R	≥32	R
Ampicillin/sulbactam	≥32	R	≥32	R	≥32	R	≥32	R	≥32	R	≥32	R
Piperacillin/tazobactam	≥128	R	≥128	R	≥128	R	≥128	R	≥128	R	≥128	R
Cefazolin	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R
Cefotetan	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R
Ceftazidime	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R
Ceftriaxone	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R
Cefepime	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R	≥64	R
Aztreonam	≥64	R	16	R	≥64	R	16	R	16	R	16	R
Ertapenem	≥8	R	≥8	R	≥8	R	≥8	R	≥8	R	≥8	R
Imipenem	≥16	R	≥16	R	≥16	R	≥16	R	≥16	R	≥16	R
Amikacin	≤2	S	≤2	S	≤2	S	≤2	S	≤2	S	≤2	S
Gentamicin	≥16	R	≤1	S	≥16	R	≤1	S	≤1	S	≤1	S
Tobramycin	≥16	R	≤1	S	≥16	R	≤1	S	≤1	S	≤1	S
Ciprofloxacin	≥4	R	≥4	R	≥4	R	≥4	R	≥4	R	≥4	R
Levofloxacin	≥8	R	≥8	R	≥8	R	≥8	R	≥8	R	≥8	R
Nitrofurantoin	≤16	S	≤16	S	≤16	S	≤16	S	≤16	S	≤16	S
Trimethoprim/ Sulfamethoxazole	≥320	R	≤20	S	≥320	R	≤20	S	≤20	S	≤20	S

Note: S=sensitive; R=resistant; MIC=minimum inhibitory concentration.

after January 26, 2023, only pDJH\_183sputum\_Plas2 was detected in isolated strains (JARVKR000000000, JARVKS000000000), indicating that pDJH\_137urine\_Plas1 was not stably maintained, while

pDJH\_183sputum\_Plas2 was (Figure 1A).

The other two plasmids, pDJH\_Plas3 (96.2 Kbp) and pDJH\_Plas4 (4.4 Kbp), do not carry any resistance genes.



B.

Characteristic	Chromosome	Plasmids of DJH_183_sputum				
		Plasmids of DJH_137 urine				pDJH_183 sputum_Plas2
		pDJH_137 urine_Plas1	pDJH_Plas3	pDJH_Plas4		
Genome size	4.8 Mb	152.2 Kbp	48.0 Kbp	96.2 Kbp	4.4 Kbp	48.6 Kbp
GC%	50.7	51.5	47	50.2	55.5	50.6
Aver CDS	909.9	716.8	640.9	680.3	426	685.3
CDS%	87.3	80.1	86.8	79.9	29.3	81.7
IS(Family)	89(20)	102(20)	32(20)	4(1)	0	34(20)









FIGURE 1. Genomic analysis and timeline of the isolation of *bla*<sub>NDM-5</sub> *E. coli* strains from the patient following COVID-19 infection — Beijing, China, 2023. (A) Isolation source and genomic composition of the strains. (B) Genomic properties of the chromosome and plasmids of the strains isolated on December 31, 2022. (C) Genomic synteny between pDJH\_137urine\_Plas1 and pDJH\_183sputum\_Plas2 plasmids. (D) Neighbor-joining tree based on the core genome of the *E. coli* strains isolated on December 31, 2022, January 5, 2023, and January 26, 2023, and *E. coli* representatives from each phylogroup. (E) Neighbor-joining tree of pDJH\_137urine\_Plas1 and the top 50 megaBLAST hits using BLAST pairwise alignments. (F) Neighbor-joining tree of pDJH\_183sputum\_Plas2 and the top 50 megaBLAST hits using BLAST pairwise alignments. (G) Neighbor-joining tree of pDJH\_NDM-5 and the top 50 megaBLAST hits using BLAST pairwise alignments. (H) Neighbor-joining tree of pDJH\_Plas3 and the top 50 megaBLAST hits using BLAST pairwise alignments. (I) Neighbor-joining tree of pDJH\_Plas4 and the top 50 megaBLAST hits using BLAST pairwise alignments.

Abbreviation: IS=Insertion Sequence; EAEC=Enteroaggregative *Escherichia coli*; ETEC=Enterotoxigenic *Escherichia coli*; EIEC=Enteroinvasive *Escherichia coli*; STEC=Shiga toxin-producing *Escherichia coli*.

## DISCUSSION

The diagnostic criteria for a UTI include clinical symptoms and the presence of a pathogen. Pyuria, or the presence of white blood cells or pus in the urine, serves as a significant indicator (9). Although the patient in this study did not report any subjective UTI

symptoms, NDM-5 *E. coli* was isolated from her urine four times over a period of 45 days, accompanied by noticeable bacteriuria and pyuria (Table 1). Consequently, a diagnosis of persistent UTI was made. The NDM-5 *E. coli* infection in this case was likely community-acquired, with the bacteria being selectively eliminated in vivo due to the patient's at-

home use of a second-generation cephalosporin. The spread of the bacterial strains appeared to be limited, as the patient's nanny did not contract the infection.

Interestingly, the NDM-5 *E. coli* was not a single clone but rather a group of homologous *E. coli* strains. These strains possessed the same chromosome but carried different plasmids and were present concurrently in multiple systems (e.g., the urethra and the respiratory tract, the urethra and the intestinal tract). Following the cessation of antibiotics, the pDJH\_137urine\_Plas1 plasmid, which conferred a broader drug resistance spectrum, gradually disappeared. In contrast, the less-drug-resistant plasmid pDJH\_183sputum\_plasmid2 remained within the NDM-5 *E. coli* population (Figure 1A).

The source of the patient's infection was investigated by collecting and analyzing their fecal samples twice after their discharge from the facility. In both instances, NDM-5 *E. coli* strains were isolated. Previous research has identified the intestinal tract as a reservoir for antibiotic-resistance genes (1,10). The patient also reported habitually wiping her perineal area from back to front after bowel movements. It was hypothesized that the *E. coli* strains' plasmids underwent recombination within the gastrointestinal tract and were subsequently transferred to the urinary tract due to the patient's wiping habit. Virulence genes, such as *fimH*, facilitated the strains' colonization along the epithelial cells of the urinary tract. Coincidentally, the bacterial strain carrying the NDM-5 plasmid, which conferred extensive antibiotic resistance, led to a persistent UTI.

In the present case, a rare occurrence of community-acquired NDM-5 *E. coli* UTI persisted for over 45 days. Two NDM-5 strains were identified, both having the same chromosome but with homologous and recombined plasmids. This finding suggests that multidrug-resistant bacteria were not stable throughout the infection. While new plasmids can be acquired, the plasmids themselves may also undergo constant recombination, leading to variations in antibiotic resistance profiles. This case implies that a diverse range of resistance genes may be intrinsic to the human body, and the CRE could undergo recombination as a consequence of environmental pressures, such as those imposed by antibiotic usage. The intestinal tract might serve as the source of the gene pool. Therefore, in contrast to exogenous infection, more attention should be given to endogenous factors contributing to the development of CRE.

**Conflicts of interest:** No conflicts of interest.

**Acknowledgements:** Charlesworth Author Services (Paper no.117766) for their critical editing and helpful

comments regarding our manuscript.

**Funding:** Supported by the National Key Research and Development Program of China (2022YFC2602203).

doi: 10.46234/ccdcw2023.110

# Corresponding authors: Xin Wang, wangxin@icdc.cn; Rongmeng Jiang, rongmengj@163.com; Biao Kan, kanbiao@icdc.cn.

<sup>1</sup> Beijing Ditan Hospital Capital Medical University, Beijing, China;

<sup>2</sup> National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China; <sup>3</sup> State Key Laboratory of Infectious Disease Prevention and Control, Beijing, China.

<sup>&</sup> Joint first authors.

Submitted: May 14, 2023; Accepted: June 20, 2023

## REFERENCES

1. Lv DY, Duan R, Fan R, Mu H, Liang JR, Xiao M, et al. *bla*<sub>NDM</sub> and *mcr-1* to *mcr-5* gene distribution characteristics in gut specimens from different regions of China. *Antibiotics (Basel)* 2021;10(3):233. <http://dx.doi.org/10.3390/antibiotics10030233>.
2. Zhou ZM, Alikhan NF, Mohamed K, Fan YL, Agama Study Group, Achtman M. The Enterobase user's guide, with case studies on *Salmonella* transmissions, *Yersinia pestis* phylogeny, and *Escherichia coli* core genomic diversity. *Genome Res* 2020;30(1):138 – 52. <http://dx.doi.org/10.1101/gr.251678.119>.
3. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al. ResFinder 4.0 for predictions of phenotypes from genotypes. *J Antimicrob Chemother* 2020;75(12):3491 – 500. <http://dx.doi.org/10.1093/jac/dkaa345>.
4. Clermont O, Christenson JK, Denamur E, Gordon DM. The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ Microbiol Rep* 2013;5(1):58 – 65. <http://dx.doi.org/10.1111/1758-2229.12019>.
5. Le Gall T, Clermont O, Gouriou S, Picard B, Nassif X, Denamur E, et al. Extraintestinal virulence is a coincidental by-product of commensalism in B2 phylogenetic group *Escherichia coli* strains. *Mol Biol Evol* 2007;24(11):2373 – 84. <http://dx.doi.org/10.1093/molbev/msm172>.
6. Brons JK, Vink SN, de Vos MGJ, Reuter S, Dobrindt U, van Elsas JD. Fast identification of *Escherichia coli* in urinary tract infections using a virulence gene based PCR approach in a novel thermal cycler. *J Microbiol Methods* 2020;169:105799. <http://dx.doi.org/10.1016/j.mimet.2019.105799>.
7. Lin WH, Wang MC, Liu PY, Chen PS, Wen LL, Teng CH, et al. *Escherichia coli* urinary tract infections: host age-related differences in bacterial virulence factors and antimicrobial susceptibility. *J Microbiol Immunol Infect* 2022;55(2):249 – 56. <http://dx.doi.org/10.1016/j.jmii.2021.04.001>.
8. Bunduki GK, Heinz E, Phiri VS, Noah P, Feasey N, Musaya J. Virulence factors and antimicrobial resistance of uropathogenic *Escherichia coli* (UPEC) isolated from urinary tract infections: a systematic review and meta-analysis. *BMC Infect Dis* 2021;21(1):753. <http://dx.doi.org/10.1186/s12879-021-06435-7>.
9. Semeniuk H, Church D. Evaluation of the leukocyte esterase and nitrite urine dipstick screening tests for detection of bacteriuria in women with suspected uncomplicated urinary tract infections. *J Clin Microbiol* 1999;37(9):3051 – 2. <http://dx.doi.org/10.1128/jcm.37.9.3051-3052.1999>.
10. Kent AG, Vill AC, Shi QJ, Satlin MJ, Brito IL. Widespread transfer of mobile antibiotic resistance genes within individual gut microbiomes revealed through bacterial Hi-C. *Nat Commun* 2020;11(1):4379. <http://dx.doi.org/10.1038/s41467-020-18164-7>.

## SUPPLEMENTARY MATERIALS

SUPPLEMENTARY TABLE S1. Specific primers for individual plasmids and shared primers for pDJH\_137urine\_Plas1 and pDJH\_183sputum\_Plas2.

Target plasmid	Primer name	5'→3' sequence	Amplicon length (bp)
pDJH_137urine_Plas1	S137P1-F1	TTTTGCCGTTACGCACCACTC	1,033
	S137P1-R1	CGCCCTGCCTCTTACTCTACT	
	S137P1-F2	AAAACGGCAAACCAGTCTCTCA	504
	S137P1-R2	CAAACGGTCTTACTCCATTCA	
pDJH_183sputum_Plas2	183_plas2F	ATGGCAAACCTGAAACGG	616
	183_plas2R	CCTGTCCCTACACCAATAA	
	183P2-5F	GGCTTACGCCACCATCAAAG	287
	183P2-5R	TACGCCAAGACGAACCATCA	
pDJH_Plas_NDM-5	S137P2-ndm-F1	GCGTAGCGTTTCCATAGCG	678
	S137P2-ndm-R1	CGAGGTCTTCGTCCTGTCC	
	S137P2-ndm-F2	CCACGACAGCCAGGAGT	522
	S137P2-ndm-R2	TTCGGGCGGTTTAGTTT	
pDJH_Plas3	S183P3-F1	AAGGCGCTTTCTGATTTGG	667
	S183P3-R1	CGCTACCTGGGTCGTTATCT	
	S183P3-F2	CCACGCTGATGGAACCTGAA	941
	S183P3-R2	GCGGAGGTCTGAATAAACACG	
pDJH_Plas4	S137P4-F1	CACATTCGCTCGCTACGCT	816
	S137P4-R1	CACGCTCACCTGCTATACGC	
	S137P4-F2	CAGTCGTTACACTACACCCA	1,175
	S137P4-R2	GGCAGGATAACCCGCTACAT	
pDJH_137urine_Plas1 pDJH_183sputum_Plas2	S183P2-F1	AGATAACGACCCAGGTAGCG	799
	S183P2-R1	AACTGAAGCGGAGATGAAGG	
	S183P2-F2	TACCGCTGTCCACGCCTTAT	598
	S183P2-R2	CACGCTACCTGGGTCGTTATCT	

SUPPLEMENTARY TABLE S2. Presence of the plasmids in *bla*<sub>NDM-5</sub> *E. coli* from a UTI patient.

Plasmid	December 31, 2022		January 5, 2023	January 26, 2023		February 17, 2023	
	Urine	Sputum	Urine	Urine	Feces	Urine	Feces
pDJH_137urine_Plas1	+	-	+	-	-	-	-
pDJH_183sputum_Plas2	-	+	-	+	+	+	+
pDJH_Plas_NDM-5	+	+	+	+	+	+	+
pDJH_Plas3	+	+	+	+	+	+	+
pDJH_Plas4	+	+	+	+	+	+	+

Note: "+" means positive; "-" means negative.