

Preplanned Studies

Occurrence and Reduction of Hepatitis B Vaccine Hesitancy Among Medical University Students — Shanxi Province, China, 2020

Huijuan Kong^{1,2,3}; Xufeng Cui^{1,2,3}; Ying Zhang^{1,2,3}; Qiang Xue¹; Jing Shi^{1,2,3}; Tian Yao^{2,4}; Yana Guo^{1,2,3}; Xiuyang Xu^{1,2,3}; Suping Wang^{1,2,3,#}; Yongliang Feng^{1,2,3,#}

Summary

What is already known about this topic?

Previous research has primarily examined the issue of hepatitis B vaccine hesitancy in migrant workers and other adult populations. However, there is a lack of studies that have specifically investigated the prevalence of hepatitis B vaccine hesitancy among university students.

What is added by this report?

In this study, 19.84% of students expressed hesitancy towards receiving the hepatitis B immunization. A negative correlation was observed between hepatitis B vaccine hesitancy and knowledge, attitudes, and practices related to hepatitis B. Conversely, a positive relationship was identified between hepatitis B-related knowledge and attitudes and practices.

What are the implications for public health practice?

This study examines the factors contributing to vaccine hesitancy towards hepatitis B at a medical university in China. The results have significant implications for developing strategies to improve hepatitis B vaccination rates.

Hepatitis B virus (HBV) infection is a significant global public health concern (1). According to the World Health Organization (WHO), in 2019, an estimated 296 million individuals worldwide had chronic hepatitis B, with 1.5 million new cases reported annually and approximately 820,000 deaths (2). In China, universal hepatitis B vaccination for newborns was introduced in 1992, resulting in a high vaccination rate among university students. However, the antibody response to HBsAg (anti-HBs) decreases over time after vaccination (3). Additionally, university students are at a higher risk of HBV infection due to various factors such as tattoos, unprotected sexual contact, pierced ears, and accidental blood exposure to

medical students. The WHO recommends booster vaccination for high-risk populations (4). However, there is a growing reluctance among individuals to receive recommended vaccinations (5). Despite this, there have been limited efforts to evaluate hepatitis B vaccine hesitancy among medical university students. This study was conducted among the class of 2017 at a medical university from April to June 2020, with a total of 1,003 students from different professions completing a questionnaire. The results illustrated that hepatitis B-related knowledge, attitudes, and practices had a negative impact on hepatitis B vaccine hesitancy. The findings from this study provide insights for developing a health education-based hepatitis B vaccination strategy.

A cross-sectional study was conducted in the class of 2017. Out of the 27 professions, nine were randomly selected for recruitment, and all students in those nine professions were included in the study. Prior to enrollment, written informed consent was obtained from all participants. A self-designed questionnaire, consisting of 15 questions on HBV knowledge, 15 questions on HBV attitudes, and 10 questions on practices (Supplementary Table S1, available at <https://weekly.chinacdc.cn/>), was distributed through the Wenjuanxing platform. The questionnaire was administered on-site by trained investigators. Students were asked to independently complete the questionnaire. The internal consistency of the questionnaire, measured by Cronbach's α , was determined to be 0.855, indicating good reliability.

Hepatitis B knowledge was categorized into four dimensions based on their relevance to infectious disease epidemics among humans: etiological knowledge (EK), transmission route knowledge (TRK), prevention measures knowledge (PMK), and epidemic status knowledge (ESK). Each correct answer was scored 1 point, while incorrect or unclear answers received 0 points. Hepatitis B attitudes were assessed

using a 5-point Likert scale, ranging from “very consistent” to “very inconsistent,” with scores of 5 to 1, respectively, indicating personal inclination. Questions 3, 4, 5, 12, 13, and 14 were reverse-scored. Hepatitis B practices were divided into two dimensions based on behavioral characteristics: health-seeking practices (HSP) and high-risk practices (HRP). A score of 1 point was given for “yes” responses to hepatitis B practice items, while “no” or “unknown” responses received 0 points. Knowledge, attitudes, and practices related to hepatitis B were classified into two groups based on the mean score.

In our study, “vaccine hesitancy” (VH) was defined as the participants’ response of “unsure”, “basically inconsistent”, or “very inconsistent” when asked about their willingness to be vaccinated against hepatitis B in the absence of detectable anti-HBs. This term was used to describe the hesitation towards receiving the hepatitis B vaccine.

The database was established using Microsoft Excel (version 2016; Microsoft Corp, Redmond, USA), and analyses were conducted using SAS (version 9.4; SAS Institute, Cary, NC, USA). The chi-square test was used to assess the difference between the groups with vaccine hesitancy and those without. Logistic regression analysis was employed to examine the associations between the independent variables and hepatitis B vaccine hesitancy. Independent variables with $P < 0.05$ in the univariate analysis were included in the logistic regression model. Structural equation modeling (SEM) was performed using Amos software (web version 24.0; IBM, New York, USA) to elucidate the interrelationships between latent variables. The model parameters were estimated using maximum likelihood estimation.

A total of 1,012 students were contacted, and 1,003 students responded, resulting in a response rate of 99%. The respondents were predominantly female (75.37%), with a sex ratio of 0.33:1. Among the respondents, the majority ($n=961$) were of Han ethnicity. The participants consisted of 747 medical professionals (74.48%) and 256 nonmedical professionals (25.52%) (Table 1).

The Kaiser-Meyer-Olkin coefficient was found to be 0.785, indicating satisfactory sampling adequacy. Additionally, Bartlett’s test showed statistical significance ($P < 0.001$), supporting the factor analysis. Based on eigenvalues, a total of four factors were extracted, explaining 69.54% of the cumulative variance. These factors were identified as attitude toward the source of infection (ASI), self-protection

attitude (SPA), life contact attitude (LCA), and vaccine effect attitude (VEA).

A total of 199 respondents expressed hesitancy towards the hepatitis B vaccine, resulting in a vaccine hesitation rate of 19.84% (199/1,003). In the univariate analysis, significant associations were observed between vaccine hesitancy and factors such as knowledge, attitudes, and health-seeking practices ($P < 0.05$) (Table 1).

The logistic regression model revealed that university students with lower TRK and PMK scores had a higher likelihood of hesitating to receive the hepatitis B vaccine ($OR=1.947$, 95% CI : 1.309, 2.897; $OR=2.501$, 95% CI : 1.682, 3.719). Participants who held stigmatizing attitudes toward the source of infection were 1.854 times more likely to hesitate in getting the hepatitis B vaccine compared to those with more inclusive attitudes (95% CI : 1.280, 2.684). Vaccine hesitancy was also associated with concerns about vaccine efficacy ($OR=6.529$, 95% CI : 4.417, 9.650). Furthermore, individuals who adopted fewer health-seeking behaviors were more likely to hesitate to receive the vaccine and had a 2.119-fold increased likelihood of adopting such behaviors (95% CI : 1.357, 3.308) (Table 2).

SEM was developed using the knowledge, attitude, and practice theory model as a basis. The initial model included knowledge-related latent variables as independent variables, attitude and behavior-related latent variables as mediators, and the observed variable VH as the dependent variable. Non-significant paths between latent variables in the hypothetical model ($P > 0.05$) were removed and corrected using the correction index. Model fit measurements indicated that the revised model demonstrated a satisfactory fit, and the model was professionally interpreted. The reference standards for the fit indicators, as well as the fit indicators for the final model, are provided in Supplementary Table S2 (available at <https://weekly.chinacdc.cn/>). A diagram of the standardized model can be seen in Figure 1. The values assigned to each variable in the revised model are presented in Supplementary Table S3 (available at <https://weekly.chinacdc.cn/>).

Supplementary Table S4 (available at <https://weekly.chinacdc.cn/>) presents the associations between different factors (TRK, PMK, ASI, HSP, LCA) and vaccine hesitancy for hepatitis B. TRK had an indirect effect on vaccine hesitancy [standard indirect effect (SIE)=-0.301]. PMK, on the other hand, influenced vaccine hesitancy through both direct and indirect

TABLE 1. Characteristics of university students with hepatitis B vaccine hesitancy [*n* (%)].

Variables	N	Hepatitis B vaccine hesitancy		Univariate analysis	
		Yes (n=199)	No (n=804)	χ^2	P value
Gender				10.935	0.001
Male	247 (24.63)	67 (27.13)	180 (72.87)		
Female	756 (75.37)	132 (17.46)	624 (82.54)		
Locality				< 0.001	0.998
Urban	368 (36.69)	73 (19.84)	295 (80.16)		
Rural	635 (63.31)	126 (19.84)	509 (80.16)		
Ethnicity				0.851	0.356
Han ethnicity	961 (95.81)	193 (20.08)	768 (79.92)		
Other	42 (4.19)	6 (14.29)	36 (85.71)		
Medical Professional				9.766	0.002
Yes	747 (74.48)	131 (17.54)	616 (82.46)		
No	256 (25.52)	68 (26.56)	188 (73.44)		
Etiological knowledge				62.620	<0.001
Good mastery (≥ 1.82)	662 (66.00)	84 (12.69)	578 (87.31)		
Poor mastery (<1.82)	341 (34.00)	115 (33.72)	226 (66.28)		
Prevention measures knowledge				68.217	<0.001
Good mastery (≥ 1.44)	605 (60.32)	69 (11.40)	536 (88.60)		
Poor mastery (<1.44)	398 (39.68)	130 (32.66)	268 (67.34)		
Transmission routes knowledge				51.415	<0.001
Good mastery (≥ 5.12)	686 (68.39)	94 (13.70)	592 (86.30)		
Poor mastery (<5.12)	317 (31.61)	105 (33.12)	212 (66.88)		
Epidemic status knowledge				31.464	<0.001
Good mastery (≥ 0.57)	570 (56.83)	78 (13.68)	492 (86.32)		
Poor mastery (<0.57)	433 (43.17)	121 (27.94)	312 (72.06)		
Attitude toward the source of infection				45.103	<0.001
Inclusion (≥ 8.94)	630 (62.81)	84 (13.33)	546 (86.67)		
Stigmatization (<8.94)	373 (37.19)	115 (30.83)	258 (69.17)		
Life contact attitude				32.020	<0.001
Inclusion (≥ 18.12)	398 (39.68)	44 (11.06)	354 (88.94)		
Stigmatization (<18.12)	605 (60.32)	155 (25.62)	450 (74.38)		
Self-protection attitude				20.715	<0.001
Inclusion (≥ 8.04)	436 (43.47)	115 (26.38)	321 (73.62)		
Stigmatization (<8.04)	567 (56.53)	84 (14.81)	483 (85.19)		
Vaccine effect attitude				104.763	<0.001
Inclusion (≥ 9.99)	551 (54.94)	45 (8.17)	506 (91.83)		
Stigmatization (<9.99)	452 (45.06)	154 (34.07)	298 (65.93)		
Health-seeking practices				18.168	<0.001
More practices (≥ 1.06)	295 (29.41)	34 (11.53)	261 (88.47)		
Fewer practices (<1.06)	708 (70.59)	165 (23.31)	543 (76.69)		
High-risk practices				0.069	0.794
More practices (≥ 1.25)	376 (37.49)	73 (19.41)	303 (80.59)		
Fewer practices (<1.25)	627 (62.51)	126 (20.10)	501 (79.90)		

TABLE 2. Multivariate analysis of hesitancy towards hepatitis B vaccine among university students.

Variables	β	S \bar{x}	Wald χ^2	OR (95% CI)*	P value
Transmission routes knowledge					
Good mastery				1	
Poor mastery	0.666	0.203	10.801	1.947 (1.309, 2.897)	0.001
Prevention measures knowledge					
Good mastery				1	
Poor mastery	0.917	0.203	20.492	2.501 (1.682, 3.719)	<0.001
Attitude toward the source of infection					
Inclusion				1	
Stigmatization	0.617	0.189	10.686	1.854 (1.280, 2.684)	0.001
Life contact attitude					
Inclusion				1	
Stigmatization	0.663	0.210	10.002	1.940 (1.287, 2.925)	0.002
Vaccine effect attitude					
Inclusion				1	
Calculation	1.876	0.199	88.573	6.529 (4.417, 9.650)	<0.001
Health-seeking practices					
More behaviors				1	
Fewer behaviors	0.751	0.227	10.913	2.119 (1.357, 3.308)	0.001

Abbreviation: OR=odds ratio; CI=confidence interval.

* Adjusted by Gender, Medical Professional, epidemic status knowledge, and self-protection attitude.




FIGURE 1. Revised model (SEM).

Note: The ellipse and arrow in the structural model represent the effects of factors. One-way straight arrows in the structural model depict the paths of each factor with standardized regression weights, and one-way straight arrows in the measurement model show the measurement loadings of each item in the scale.

Abbreviation: PMK=prevention measures knowledge; ASI=attitude toward the source of infection; PMK=prevention measures knowledge; VH=vaccine hesitancy; LCA=life contact attitude; TRK=transmission route knowledge; HSP=health-seeking practices.

effects (standard direct effect (SDE)=-0.168, SIE=-0.101). ASI (SDE=-0.267) and HSP (SDE=-0.204) only had direct effects on hepatitis B vaccine hesitancy. Additionally, LCA had an indirect effect on vaccine hesitancy (SIE=-0.066), and this relationship was partially mediated by ASI. The analysis also revealed a negative correlation between hepatitis B-related knowledge, attitudes, and practices, and vaccine hesitancy. Further details regarding the structural model parameters can be found in [Supplementary Table S5](#) (available at <https://weekly.chinacdc.cn/>). These findings indicate the importance of health education that focuses on enhancing knowledge, attitudes, and practices related to hepatitis B, as part of an effective hepatitis B vaccination strategy.

DISCUSSION

The survey results revealed that the rate of vaccine hesitancy for hepatitis B among university students was 19.84%, which was lower compared to other migrant workers (6). This difference may be attributed to the medical students' curriculum, specifically their education on infectious diseases, which contributes to a more accurate comprehension of hepatitis B-related knowledge.

Previous research has indicated that being aware of vaccine-preventable diseases can encourage vaccination (7). Our study found that students who were hesitant about vaccines had significantly less knowledge about hepatitis B compared to students who were accepting of vaccines. SEM also showed that knowledge had an indirect impact on vaccine hesitancy. These results are consistent with previous studies, highlighting the importance of increasing awareness of hepatitis B-related knowledge among university students. This can be achieved through initiatives like campus-wide events and interdisciplinary curricula, which can help mitigate vaccine hesitancy.

Stigma presents a significant obstacle to the effective implementation of preventive measures against HBV infection (8). Our study found that students who expressed stigmatizing attitudes towards hepatitis B patients and carriers were more likely to hesitate in getting vaccinated against hepatitis B. These findings highlight the need for universities and government organizations to provide early and positive guidance to enhance student education and promote understanding of the transmission routes of HBV and other viruses, in order to reduce counterproductive stigmatization among university students.

A survey conducted by the WHO and UNICEF from 2014 to 2016 revealed that doubts regarding vaccine safety and concerns about potential side effects were significant factors contributing to vaccine hesitancy (9). Our study, consistent with previous research on vaccine hesitancy in other contexts (10), identified concerns about the effectiveness and safety of vaccines as crucial determinants for refusing vaccination or lacking the intention to receive it ($OR=6.529$). We observed that university students who exhibited worry towards the hepatitis B vaccine were more likely to demonstrate vaccine hesitancy, which can have a negative impact on vaccine confidence and communication with both their families and future patients. Thus, our findings underscore the importance of educating university students about the safety and effectiveness of vaccines, as well as raising awareness about the critical role vaccines play in disease prevention. These ongoing issues persistently contribute to confusion and uncertainty surrounding vaccines.

Risk factors associated with hepatitis B vaccine hesitancy among university students include inadequate knowledge, stigmatization of attitudes towards the source of hepatitis B infection, and lacking health-seeking behaviors. Poor understanding, unfavorable attitudes, and inadequate practices related to hepatitis B are associated with increased vaccine hesitancy. Therefore, prioritizing knowledge dissemination is crucial for the development of effective hepatitis B vaccination strategies. Implementing educational programs at universities plays a vital role in raising awareness about prevention and enhancing vaccine coverage among students.

Previous research has primarily examined hepatitis B vaccine hesitancy in migrant workers, with limited attention given to medical university students. However, understanding the factors influencing vaccine hesitancy among medical students and developing effective health education strategies is crucial. This study aimed to address this gap by evaluating hepatitis B vaccine hesitancy in a medical university setting.

This study was subject to some limitations. First, the study utilized closed-ended questions, which limited the students' ability to provide nuanced insights into their knowledge, attitudes, and practices regarding hepatitis B. Second, this study was conducted in a single medical university, and therefore may not fully capture the current status and determinants of hepatitis B vaccine hesitancy among Chinese university

students. Future surveys should include a more diverse range of universities to ensure a representative sample.

Conflicts of interest: No conflicts of interest.

doi: [10.46234/ccdcw2024.081](https://doi.org/10.46234/ccdcw2024.081)

Corresponding authors: Yongliang Feng, yongliang.feng@sxmu.edu.cn; Suping Wang, supingwang@sxmu.edu.cn.

¹ School of Public Health, Shanxi Medical University, Taiyuan City, Shanxi Province, China; ² Center of Clinical Epidemiology and Evidence Based Medicine, Shanxi Medical University, Taiyuan City, Shanxi Province, China; ³ Key Laboratory of Coal Environmental Pathogenicity and Prevention (Shanxi Medical University), Ministry of Education, Taiyuan City, Shanxi Province, China; ⁴ First Hospital/First Clinical Medical College of Shanxi Medical University, Taiyuan City, Shanxi Province, China.

Submitted: December 15, 2023; Accepted: January 12, 2024

REFERENCES

- Chen SQ, Li J, Wang D, Fung H, Wong LY, Zhao L. The hepatitis B epidemic in China should receive more attention. *Lancet* 2018;391(10130):1572. [https://doi.org/10.1016/S0140-6736\(18\)30499-9](https://doi.org/10.1016/S0140-6736(18)30499-9).
- World Health Organization. Hepatitis B. 2023. <https://www.who.int/news-room/fact-sheets/detail/hepatitis-b>. [2023-9-8].
- Alshehri AA, Alshamrani SA, Elnoubi OA, Alshahrani AJ, Saif AM, Nahari MH, et al. Persistence of hepatitis B surface antibody to hepatitis B vaccine among medical college students in Najran University, Saudi Arabia. *Bioinformation* 2022;18(7):617 – 21. <https://doi.org/10.6026/97320630018617>.
- World Health Organization. Hepatitis B vaccines: WHO position paper, July 2017 - Recommendations. *Vaccine* 2019;37(2):223 – 5. <https://doi.org/10.1016/j.vaccine.2017.07.046>.
- Gilbert-Esparza E, Brady A, Haas S, Wittstruck H, Miller J, Kang Q, et al. Vaccine hesitancy in college students. *Vaccines* 2023;11(7):1243. <https://doi.org/10.3390/vaccines11071243>.
- Liu C, Nicholas S, Wang J. The association between protection motivation and hepatitis b vaccination intention among migrant workers in Tianjin, China: a cross-sectional study. *BMC Public Health* 2020;20(1):1219. <https://doi.org/10.1186/s12889-020-09292-2>.
- Pires C. Global predictors of COVID-19 vaccine hesitancy: a systematic review. *Vaccines* 2022;10(8):1349. <https://doi.org/10.3390/vaccines10081349>.
- Nguyen TTL, Pham TTH, So S, Hoang THV, Nguyen TTU, Ngo TB, et al. Knowledge, attitudes and practices toward hepatitis B virus infection among students of medicine in Vietnam. *Int J Environ Res Public Health* 2021;18(13):7081. <https://doi.org/10.3390/ijerph18137081>.
- Lane S, MacDonald NE, Marti M, Dumolard L. Vaccine hesitancy around the globe: analysis of three years of WHO/UNICEF Joint Reporting Form data-2015–2017. *Vaccine* 2018;36(26):3861 – 7. <https://doi.org/10.1016/j.vaccine.2018.03.063>.
- Šálek J, Čelko AM, Dáňová J. Vaccination perception and attitude among undergraduate medical and teacher education students at Charles university, Prague, Czech republic. *Vaccines* 2020;8(1):136. <https://doi.org/10.3390/vaccines8010136>.

SUPPLEMENTARY MATERIAL

SUPPLEMENTARY TABLE S1. Knowledge, attitudes, and practices regarding Hepatitis B among university students.

Latent variables	Items
Etiological knowledge	<ol style="list-style-type: none"> 1. Positive hepatitis B surface antigen (HBsAg) indicates HBV infection. 2. Hepatitis B surface antibody (anti-HBs) is a protective antibody. 3. Hepatitis B virus carriers are the source of hepatitis B infection.
Transmission routes knowledge	<ol style="list-style-type: none"> 4. HBV can be transmitted through broken skin or mucous membranes. 5. HBV can be transmitted by blood-sucking insects. 6. HBV can be transmitted through pedicures, tattoos, and earrings. 7. HBV can be transmitted by sharing syringes, razors, and dental tools. 8. HBV can be transmitted through blood or blood products. 9. HBV can be transmitted by droplets and dust. 10. HBV can be transmitted through sexual intercourse. 11. HBV can be transmitted from mother to child. 12. HBV can be transmitted through daily study, work, or life contact.
Prevention measures knowledge	<ol style="list-style-type: none"> 13. Hepatitis B vaccination is the primary method of preventing hepatitis B virus infection. 14. Hepatitis B vaccination is recommended when Hepatitis B Antibody is negative.
Epidemic status knowledge	<ol style="list-style-type: none"> 15. China is now a medium endemic country for hepatitis B virus infection.
Attitude toward the source of infection	<ol style="list-style-type: none"> 1. Hepatitis B patients' privacy should be protected. 2. Hepatitis B patients should not be discriminated against.
Self-protection attitude	<ol style="list-style-type: none"> 3. Fear that you may be infected with the hepatitis B virus at any time. 4. Fear of others knowing you are a hepatitis B carrier or patient. 5. Living in the same dormitory with hepatitis B patients can be psychologically stressful.
Life contact attitude	<ol style="list-style-type: none"> 6. Willing to be friends with chronic asymptomatic HBV carriers. 7. Willingness to share meals with chronic asymptomatic HBV carriers. 8. Willingness to live with chronic asymptomatic HBV carriers. 9. Willingness to have close contact with the hepatitis B virus carriers. 10. Willingness to marry hepatitis B virus carriers. 11. Willingness to treat people with hepatitis B as others.
Vaccine effect attitude	<ol style="list-style-type: none"> 12. Fear of side effects of hepatitis B vaccine. 13. Fear that the hepatitis B vaccine does not provide enough protection against the hepatitis B virus. 14. Worried that my knowledge of the hepatitis B vaccine is not very comprehensive.
Health-seeking practices	<ol style="list-style-type: none"> 1. Hepatitis B vaccination other than birth. 2. Tested for HBV serological markers and/or hepatitis B virus DNA in the last three years. 3. Will take the initiative to pay attention to the knowledge about hepatitis B.
High-risk practices	<ol style="list-style-type: none"> 4. Pedicure. 5. Tattoo, eyebrow tattoo. 6. Manicure. 7. Pierced ears. 8. Shared eyebrow trimmer, shaver. 9. Shared toothbrush. 10. Having sexual intercourse.

SUPPLEMENTARY TABLE S2. Goodness of fit indices of structural equation model.

Indicators	GFI	AGFI	NFI	IFI	CFI	RMR	RMSEA
Reference standards	>0.90	>0.90	>0.90	>0.90	>0.90	<0.05	<0.05
Model results	0.972	0.960	0.971	0.982	0.982	0.024	0.040

Abbreviation: GFI=goodness fit index; AGFI=adjusted goodness of fit index; NFI=normed fit index; IFI=incremental fit index; CFI=comparative fit index; RMR=root mean square residual; RMSEA=root mean square error of approximation.

SUPPLEMENTARY TABLE S3. Revised model (SEM) variables related to hepatitis B vaccine hesitancy among university students.

Latent variables	Items	Assignment
TRK	Q1 HBV can be transmitted through broken skin or mucous membranes	1=correct, 2=wrong, 3=unclear
	Q2 HBV can be transmitted by blood-sucking insects	1=correct, 2=wrong, 3=unclear
	Q3 HBV can be transmitted through pedicures, tattoos, and earrings	1=correct, 2=wrong, 3=unclear
	Q4 HBV can be transmitted through blood or blood products	1=correct, 2=wrong, 3=unclear
	Q5 HBV can be transmitted through sexual intercourse	1=correct, 2=wrong, 3=unclear
PMK	Q6 Hepatitis B vaccination is the primary method of preventing hepatitis B virus infection	1=correct, 2=wrong, 3=unclear
	Q7 Hepatitis B vaccination is recommended when Hepatitis B antibody is negative	1=correct, 2=wrong, 3=unclear
LCA	Q8 Willing to be friends with chronic asymptomatic HBV carriers	1=Fully conform, 2=Basically conform, 3=Uncertain, 4=Basically not conform, 5=Not at all conform
	Q9 Willingness to share meals with chronic asymptomatic HBV carriers	1=Fully conform, 2=Basically conform, 3=Uncertain, 4=Basically not conform, 5=Not at all conform
	Q10 Willingness to live with chronic asymptomatic HBV carriers	1=Fully conform, 2=Basically conform, 3=Uncertain, 4=Basically not conform, 5=Not at all conform
ASI	Q11 Hepatitis B patients' privacy should be protected.	1=Fully conform, 2=Basically conform, 3=Uncertain, 4=Basically not conform, 5=Not at all conform
	Q12 Hepatitis B patients should not be discriminated against	1=Fully conform, 2=Basically conform, 3=Uncertain, 4=Basically not conform, 5=Not at all conform
VH	Q13 Detected no hepatitis B antibody and willing to receive hepatitis B vaccination	1=Fully conform, 2=Basically conform, 3=Uncertain, 4=Basically not conform, 5=Not at all conform
HSP	Q14 Hepatitis B vaccination other than birth	1=Yes, 2=No, 3=Unknown
	Q15 Tested for HBV serological markers and/or hepatitis B virus DNA in the last three years	1=Yes, 2=No, 3=Unknown

Abbreviation: TRK=transmission route knowledge; PMK=prevention measures knowledge; LCA=life contact attitude; ASI=attitude toward the source of infection; VH=vaccine hesitancy; HSP=health-seeking practices.

SUPPLEMENTARY TABLE S4. Effect analysis between the latent variable and hepatitis B vaccine hesitancy status.

Latent variable	Hepatitis B vaccine hesitancy		
	Direct effect	Indirect effect	Total effect
Transmission routes knowledge	–	–0.301	–0.301
Preventive measures Knowledge	–0.168	–0.101	–0.296
Attitude toward the source of infection	–0.267	–	–0.267
Life contact attitude	–	–0.066	–0.066
health-seeking practices	–0.204	–	–0.204

Note: “–” means no corresponding effect.

SUPPLEMENTARY TABLE S5. Parameters of the structural model in the university.

Path	Unstandardized coefficients	Standardized coefficients	CR	P value
Hepatitis B vaccine hesitancy – prevention measures knowledge	-0.180	-0.168	-4.531	***
Hepatitis B vaccine hesitancy – attitude towards the source of infection	-0.291	-0.267	-7.696	***
Hepatitis B vaccine hesitancy – health-seeking practices	-0.370	-0.204	-4.843	***
Life contact attitude – prevention measures knowledge	0.174	0.187	5.301	***
Attitude to the source of infection – prevention measures knowledge	0.327	0.332	8.889	***
Attitude towards the source of infection – life contact attitude	0.264	0.249	6.940	***
Prevention measures knowledge – transmission routes knowledge	0.928	0.901	27.857	***
Health-seeking practices – transmission routes knowledge	0.178	0.292	4.634	***

*** $P < 0.001$.