

## Preplanned Studies

## The Association Between Multidimensional Influencing Factors and Depressive Symptoms in Chinese Adolescents — Beijing Municipality, China, July 2022–April 2023

Jia Zhou<sup>1,2</sup>; Hai Zhao<sup>3</sup>; Ruilan Zhao<sup>4</sup>; Zizhao Feng<sup>1</sup>; Bin Dong<sup>2,#</sup>

### Summary

#### What is already known about this topic?

Depression significantly impacts the mental health of Chinese adolescents. Identifying risk factors specific to adolescent depression is crucial for prioritizing intervention strategies.

#### What is added by this report?

Neuroticism and emotional abuse were associated with an increased risk of depressive symptoms, whereas a positive coping style was directly and strongly associated with a decreased risk.

#### What are the implications for public health practice?

Parental awareness of emotional abuse is critical in addressing adolescent depression. Future intervention strategies should aim to enhance individuals' positive coping mechanisms to improve mental health outcomes.

Depressive symptoms significantly affect the mental health of Chinese adolescents (1), with previous research indicating a prevalence rate of 24.3% (2). The presence of depressive symptoms in this demographic is particularly concerning due to its established association with several adverse outcomes. These include extensive and prolonged impairments in interpersonal relations, social interaction, educational achievement, and work performance (3). Moreover, depressive symptoms are linked to substance misuse and an increased risk of suicide (4). Although the precise causes of depression remain unclear, it is crucial to identify risk factors that are unique to adolescent depression to better prioritize intervention strategies. Bronfenbrenner's ecological systems theory emphasizes that both individual characteristics and various environmental levels influence child and adolescent development (5). Despite previous studies investigating individual factors of adolescent depression, there remains a deficiency in comprehensive models that

consider multidimensional influences and the interactions among different risk factors.

To thoroughly examine the factors influencing depressive symptoms in adolescents, identify the most significant ones, and delineate prevention targets, we conducted a cross-sectional study in eight secondary schools in Beijing. The study aimed to analyze the association between depressive symptoms and various factors such as personality, coping style, childhood abuse, school bullying, parent and peer attachment, and perceived stress among Chinese adolescents. Data collection occurred from July 7, 2022, to April 9, 2023, employing convenience sampling through a mix of online and field surveys. Rigorous data verification and quality control measures were implemented to ensure the integrity and accuracy of the data. Quality control personnel were assigned to verify the logic and completeness of the responses, achieving a questionnaire completion rate of 92.8% (1,671/1,800). The study protocol received approval from the Ethics Committee of Beijing Anding Hospital, and written informed consent was obtained from all participants and their parents.

We administered the 20-item Center for Epidemiologic Studies Depression Scale for Children (CES-DC), a self-report screening tool designed to assess depressive symptoms in adolescents. Scores exceeding 20 are indicative of depressive symptoms in the general population. On an individual level, personality and coping styles were evaluated. Personality was assessed using the Revised Eysenck Personality Questionnaire (EPQ-R), which measures four traits across 88 items: extraversion, neuroticism, psychoticism, and a lie scale, with higher scores suggesting a greater degree of the respective personality traits. Coping styles were assessed using the Simplified Coping Style Questionnaire (SCSQ), which consists of two dimensions: positive coping (12 items) and negative coping (8 items), where higher scores on each dimension indicate more pronounced coping strategies.

At the level of school and family environments, factors such as childhood abuse and neglect, parental and peer attachments, and peer victimization were considered. Childhood abuse and neglect were quantified using the Childhood Trauma Questionnaire-Short Form (CTQ-SF), which comprises 28 items across five dimensions: emotional abuse, physical abuse, sexual abuse, emotional neglect, and physical neglect. Higher scores on each dimension denote a higher level of reported abuse or neglect. Parental and peer attachments were assessed using the Inventory of Parent and Peer Attachment (IPPA), which measures the quality of attachments to fathers and mothers separately, focusing on key aspects of attachment such as trust, communication, and alienation, with higher scores indicating a more secure parent-child relationship. Peer victimization was measured using the Multidimensional Peer-Victimization Scale (MPVS), where higher scores indicate greater victimization.

At the community and societal level, perceived stress was measured using the Chinese Perceived Stress Scale (CPSS) (6), where higher scores suggest increased perceived stress (Supplementary Table S1, available at <https://weekly.chinacdc.cn/>). All scales employed in this study demonstrate robust reliability and validity and are extensively utilized within our national context.

Descriptive data are presented as medians with interquartile ranges (IQR). The Wilcoxon rank test was employed for univariate analysis to evaluate differences between groups. Multiple linear regression analysis was then conducted to further investigate these factors and assess their collective impact. A network model was developed to estimate the relationships between depressive symptoms and associated factors using the R-package qgraph. In this model, nodes represent depressive symptoms and factors, while edges represent partial correlation coefficients between pairs of nodes. Thicker and more saturated edges indicate stronger correlations. Green edges signify positive partial correlations, and red edges indicate negative correlations. Statistical analyses were performed using SAS (version 9.4; SAS Institute, Cary, NC, US) and R Studio software (version 4.2.1; R Foundation for Statistical Computing, Vienna, Austria). All tests were two-sided and conducted at the 0.05 level of significance.

This study enrolled 1,800 adolescents, out of which 129 were excluded due to missing critical data, such as

scores on the CES-DC scale. Consequently, the analysis incorporated 1,671 participants, consisting of 810 boys and 861 girls. The mean age of the adolescents was  $14.14 \pm 1.62$  years. Participants were almost evenly split between educational levels, with 861 (51.53%) attending junior high and 810 (48.47%) attending senior high school. The average CES-DC scale score was  $14.88 \pm 10.48$ . Within the population, 444 individuals (26.57%) scored above 20 on the CES-DC scale, indicative of more significant depressive symptoms, whereas 1,227 (73.43%) scored 20 or below.

The results of the univariate analysis revealed significant distinctions in personality dimensions among adolescents with depressive symptoms. Specifically, these adolescents exhibited notably higher scores in neuroticism and psychoticism, and lower scores in extraversion and the lie dimension ( $P$  value  $< 0.001$ ). Furthermore, coping styles differed significantly; adolescents with depressive symptoms demonstrated lower scores in positive coping styles and higher scores in negative coping styles ( $P$  value  $< 0.001$ ). Environmental factors related to family and school also varied. Adolescents with depressive symptoms reported lower scores in parent-child attachment and higher scores in peer victimization and childhood abuse ( $P$  value  $< 0.001$ ). Additionally, at the mezzo level, the perceived stress among these adolescents was significantly elevated ( $P$  value  $< 0.001$ ) (Table 1). In the multiple linear regression analysis, which yielded an  $F$ -value of 92.32 and accounted for 61% of the variance ( $R^2=0.61$ ,  $P$  value  $< 0.001$ ), the retained variables included extraversion, neuroticism, positive coping, negative coping, parent-child attachment, peer victimization, emotional abuse, emotional neglect, and perceived stress (Table 2). Gender-based stratification analyses were also conducted (Supplementary Table S2, available at <https://weekly.chinacdc.cn/>).

The network analysis revealed that among the factors influencing depressive symptoms, neuroticism ( $r=0.24$ ), emotional abuse ( $r=0.22$ ), and positive coping style ( $r=-0.23$ ) exhibited the strongest and most direct correlations ( $P$  value  $< 0.001$ ). Stress demonstrated a positive correlation with neuroticism and a negative correlation with extraversion. The impact of stress on depressive symptoms appears to be predominantly mediated by personality traits, particularly neuroticism and extraversion, as shown in Figure 1.

TABLE 1. The score distribution of the influencing factors between depressive symptom groups in Chinese adolescents, median (IQR).

Variables	Total	CES-DC≤20	CES-DC>20	Z	P value
Age	14 (13–16)	13 (13–16)	15 (13–16)	3.29	0.001
Sex, n (%)				29.76	<0.001
Boy	810 (48.47)	644 (79.51)	166 (20.49)		
Girl	861 (51.53)	583 (67.71)	278 (32.29)		
Individual factors					
Personality					
Psychoticism	2 (1–4)	2 (1–3)	3 (2–5)	7.37	<0.001
Extraversion	15 (10–17)	15 (11–17)	13 (8–16)	–5.62	<0.001
Neuroticism	6 (2–14)	4 (1–9)	15 (8–19)	12.62	<0.001
Lie	16 (12–19)	16 (13–19)	14 (11–17)	–7.24	<0.001
Coping style					
Positive coping	23 (17–28)	24 (20–30)	18 (13–23)	–14.05	<0.001
Negative coping	11 (8–15)	10 (8–14)	13 (10–16)	7.59	<0.001
Family and school environment					
Parent-child attachment	77 (64–90)	81 (68–92)	66 (56–76)	–11.67	<0.001
Peer-victimization	4 (0–12)	2 (0–9)	10 (2–26)	8.94	<0.001
Childhood abuse					
Emotional abuse	6 (5–7)	5 (5–6)	8 (6–11)	17.33	<0.001
Physical abuse	5 (5–5)	5 (5–5)	5 (5–7)	11.53	<0.001
Sexual abuse	5 (5–5)	5 (5–5)	5 (5–5)	6.67	<0.001
Emotional neglect	8 (5–12)	6 (5–9)	12 (8–16)	15.26	<0.001
Physical neglect	6 (5–9)	5 (5–9)	8 (6–10)	11.40	<0.001
Total score	31 (26–39)	29 (26–34)	40 (33–48)	16.77	<0.001
Mezzo level					
Perceived stress	16 (11–20)	14 (9–18)	20 (16–24)	12.41	<0.001

Abbreviation: IQR=interquartile range; CES-DC=center for epidemiologic studies depression scale for children.

## DISCUSSION

This study examines the impact of multi-dimensional factors on depressive symptoms in adolescents, guided by ecological systems theory and network structure analysis. Findings suggest that the most direct and potent influences on depressive symptoms occur at the individual level, particularly through traits such as neuroticism and positive coping styles. Furthermore, the relationship between stress and depressive symptoms appears to be primarily moderated by personality traits, notably neuroticism and extraversion.

Our findings indicate that high neuroticism is associated with an increased risk of depressive symptoms in adolescents, while extraversion appears to serve as a protective factor against depression. Furthermore, the network analysis revealed that

neuroticism and extraversion inversely modulate the effects of perceived stress. Neuroticism, characterized by heightened negative emotionality and stress reactivity, may enhance stress appraisals and psychological distress, thereby increasing susceptibility to depressive symptoms (7). Conversely, extraversion, which is marked by sociability, assertiveness, and considerable emotional expressiveness, is typically linked with lower stress appraisals and reduced reactivity to stressful events, providing a protective effect against depressive symptoms (8). However, given that personality traits are stable and enduring, altering them to prevent depressive symptoms in adolescents is not a viable approach.

Our research has identified an additional individual factor, the positive coping style, which exerts a significant and direct impact on depressive symptoms among adolescents. A positive coping style entails

TABLE 2. Multiple linear regression analysis for the association between depressive symptoms and multi-dimensional influencing factors among Chinese adolescents.

Variables	Parameter estimation	Standard error	t	P value	VIF
Intercept	-1.91	3.33	-0.57	0.566	0
Age	0.72	0.19	3.80	<0.001	1.07
Extraversion	-0.28	0.07	-3.88	<0.001	1.23
Neuroticism	0.39	0.07	5.98	<0.001	2.22
Positive Coping	-0.30	0.05	-6.26	<0.001	1.46
Negative Coping	0.29	0.07	4.13	<0.001	1.39
Peer-Victimization	0.12	0.02	5.82	<0.001	1.19
Emotional Abuse	0.72	0.12	6.16	<0.001	1.48
Emotional Neglect	0.23	0.07	3.13	0.002	1.41
Perceived Stress	0.19	0.06	3.33	0.001	2.29

$R^2=0.61$ ,  $F=92.32$ ,  $P$  value<0.001

Note: The assumptions required for linear regression were met.

Abbreviation: VIF=variance inflation factor.




FIGURE 1. Network structure of associations between multidimensional influencing factors and depressive symptoms in Chinese adolescents.

Note: Network structures were automatically generated using the “qgraph” package in R. These networks visually represent nodes (which include depressive symptoms and influential factors) and edges (representing pairwise partial correlations among items). In the network, green edges denote positive associations, and red edges denote negative connections. Additionally, the thickness of the lines corresponds to the strength of the associations.

Abbreviation: AGE=age; DPS=depressive symptoms; PES=perceived stress; EXT=extraversion; NEU=neuroticism; PCS=positive coping style; NCS=negative coping style; EMA=emotional abuse; EMN=emotional neglect; PCA=parent-child attachment; PEV=peer-victimization.

addressing problems in a direct and rational manner and approaching challenges with an optimistic attitude. This approach can facilitate a sense of accomplishment, thereby promoting psychological health and reducing the prevalence of depressive symptoms. Our findings align with previous research,

which has reported that a positive coping style is a protective factor against depression (9). Conversely, a negative coping style, which is characterized by avoidance, withdrawal, and denial, is associated with an increased risk of depressive symptoms. Unlike inherent personality traits, a positive coping style can

be developed through education and cognitive behavioral therapy, making it a crucial target for intervention.

Consistent with prior research, our findings confirm that emotional abuse constitutes a particularly detrimental form of childhood maltreatment (10) for the development of depressive symptoms in adolescents. It is hypothesized that exposure to emotional abuse, characterized by statements such as “you are such a stupid child; you are worthless,” can evoke feelings of assault, rejection, and degradation. Accordingly, affected individuals are more likely to develop negative self-perceptions and a pessimistic attributional style, which are known contributors to the onset of depression (11). However, emotional abuse has historically received less attention compared to other types of abuse, such as sexual and physical abuse. This oversight may be attributed to the chronic nature of the damage caused by emotional abuse, which is less immediately apparent and whose impacts are often underestimated. Consequently, there is a pressing need to allocate greater resources and focus toward the prevention of this form of abuse in children and adolescents.

This study is subject to some limitations. First, due to its cross-sectional design, it is impossible to establish causal relationships, and the estimations of network structure should be considered tentative and exploratory. The hypotheses generated by this study require further validation through prospective designs. Second, the sample was recruited exclusively from Beijing, which may limit the generalizability and representativeness of the findings. Future studies with larger, more nationally representative samples are needed to deepen the understanding of these results. Despite these limitations, this study is the first to depict the interrelationships among ecological systems theory-based multi-dimensional factors influencing adolescent depression using network structure. The findings provide valuable insights, offering clearer evidence for the development of more targeted interventions in the future.

In conclusion, a positive coping style has been identified as a critical protective factor against depressive symptoms in adolescents, while neuroticism and emotional abuse were notable risk factors. It is imperative to emphasize parental awareness regarding emotional abuse. Future intervention strategies should prioritize enhancing positive coping mechanisms and mitigating negative ones to bolster adolescents' mental health resilience.

**Conflicts of interest:** No conflicts of interest.

**Funding:** Supported by the Capital's Funds for Health Improvement and Research (Grant No. 2020-2-1171) and the Beijing Hospitals Authority Youth Programme (Grant No. QML20231903).

doi: [10.46234/ccdcw2024.160](https://doi.org/10.46234/ccdcw2024.160)

# Corresponding author: Bin Dong, [bindong@bjmu.edu.cn](mailto:bindong@bjmu.edu.cn).

<sup>1</sup> The National Clinical Research Center for Mental Disorders & Beijing Key Laboratory of Mental Disorders, Beijing Anding Hospital & the Advanced Innovation Center for Human Brain Protection, Capital Medical University, Beijing, China; <sup>2</sup> Institute of Child and Adolescent Health, School of Public Health, Peking University, Beijing, China; <sup>3</sup> Beijing Center for Disease Prevention and Control, Beijing, China; <sup>4</sup> Beijing Shunyi District Center for Disease Control and Prevention, Beijing, China.

Submitted: March 01, 2024; Accepted: May 09, 2024

## REFERENCES

- Li JY, Li J, Liang JH, Qian S, Jia RX, Wang YQ, et al. Depressive symptoms among children and adolescents in China: a systematic review and meta-analysis. *Med Sci Monit* 2019;25:7459 – 70. <https://doi.org/10.12659/MSM.916774>.
- Tang XF, Tang SQ, Ren ZH, Wong DFK. Prevalence of depressive symptoms among adolescents in secondary school in mainland China: a systematic review and meta-analysis. *J Affect Disord* 2019;245:498 – 507. <https://doi.org/10.1016/j.jad.2018.11.043>.
- Thapar A, Eyre O, Patel V, Brent D. Depression in young people. *Lancet* 2022;400(10352):617 – 31. [https://doi.org/10.1016/S0140-6736\(22\)01012-1](https://doi.org/10.1016/S0140-6736(22)01012-1).
- Miller L, Campo JV. Depression in adolescents. *N Engl J Med* 2021;385(5):445 – 9. <https://doi.org/10.1056/NEJMra2033475>.
- Bronfenbrenner U. Ecological systems theory. In: Kazdin AE, editor. *Encyclopedia of psychology*. Washington: American Psychological Association. 2020; p. 129-33. <https://psycnet.apa.org/record/2004-12701-046>.
- Lu W, Bian Q, Wang WZ, Wu XL, Wang Z, Zhao M. Chinese version of the perceived stress scale-10: a psychometric study in Chinese university students. *PLoS One* 2017;12(12):e0189543. <https://doi.org/10.1371/journal.pone.0189543>.
- Banjongrewadee M, Wongpakaran N, Wongpakaran T, Pipanmekaporn T, Punjasawadwong Y, Mueankwan S. The role of perceived stress and cognitive function on the relationship between neuroticism and depression among the elderly: a structural equation model approach. *BMC Psychiatry* 2020;20(1):25. <https://doi.org/10.1186/s12888-020-2440-9>.
- Olawa BD, Idemudia ES. The extraversion-neuroticism and geriatric depression relations: do social engagements and social supports have roles to play? *Heliyon* 2020;6(12):e05719. <http://dx.doi.org/10.1016/j.heliyon.2020.e05719>.
- Sun L, Zhang Y, He JY, Qiao KX, Wang C, Zhao S, et al. Relationship between psychological capital and depression in Chinese physicians: the mediating role of organizational commitment and coping style. *Front Psychol* 2022;13:904447. <https://doi.org/10.3389/fpsyg.2022.904447>.
- Zhou J, Fan AYY, Zhou XY, Pao C, Xiao L, Feng Y, et al. Interrelationships between childhood maltreatment, depressive symptoms, functional impairment, and quality of life in patients with major depressive disorder: a network analysis approach. *Child Abuse Negl* 2022;132:105787. <https://doi.org/10.1016/j.chiabu.2022.105787>.
- van Harmelen AL, de Jong PJ, Glashouwer KA, Spinhoven P, Penninx BWJH, Elzinga BM. Child abuse and negative explicit and automatic self-associations: the cognitive scars of emotional maltreatment. *Behav Res Ther* 2010;48(6):486 – 94. <https://doi.org/10.1016/j.brat.2010.02.003>.

## SUPPLEMENTARY MATERIALS

SUPPLEMENTARY TABLE S1. Chinese perceived stress scale.

No.	Items	Scale measurement				
		Never	Almost never	Sometimes	Fairly often	Very often
1	In the last month, how often have you been upset because of something that happened unexpectedly					
2	In the last month, how often have you felt that you were unable to control the important things in your life					
3	In the last month, how often have you felt nervous and "stressed"					
4	In the last month, how often have you felt confident about your ability to handle your personal problems					
5	In the last month, how often have you felt that things were going your way					
6	In the last month, how often have you found that you could not cope with all the things that you had to do					
7	In the last month, how often have you been able to control irritations in your life					
8	In the last month, how often have you felt that you were in control of things					
9	In the last month, how often have you been angered because of things that were outside your control					
10	In the last month, how often have you felt difficulties were piling up so high that you could not overcome them					

SUPPLEMENTARY TABLE S2. Multiple linear regression analysis for the association between depressive symptoms and multi-dimensional influencing factors among boys and girls.

Gender	Variables	Parameter estimation	Standard error	t	P value	VIF
Boys	Intercept	-0.59874	5.00557	-0.12	0.9049	0
	Age	0.64877	0.28248	2.30	0.0225	1.12710
	Extraversion	-0.37153	0.10462	-3.55	0.0005	1.32335
	Neuroticism	0.47863	0.09590	4.99	<0.0001	2.14730
	Positive coping	-0.29759	0.06445	-4.62	<0.0001	1.41458
	Negative coping	0.32764	0.10190	3.22	0.0015	1.39711
	Peer-victimization	0.03076	0.03564	0.86	0.3890	1.19437
	Emotional abuse	0.92251	0.18001	5.12	<0.0001	1.33073
	Emotional neglect	0.14420	0.09318	1.55	0.1230	1.28041
Perceived stress	0.17119	0.09032	1.90	0.0592	2.35108	
Girls	Intercept	-2.07750	4.27762	-0.49	0.6275	0
	Age	0.66544	0.24443	2.72	0.0068	1.04921
	Extraversion	-0.15205	0.09251	-1.64	0.1013	1.23781
	Neuroticism	0.25686	0.08213	3.13	0.0019	2.28974
	Positive coping	-0.33652	0.06761	-4.98	<0.0001	1.58220
	Negative coping	0.26141	0.08850	2.95	0.0034	1.38900
	Peer-victimization	0.15421	0.02311	6.67	<0.0001	1.19822
	Emotional abuse	0.56503	0.13808	4.09	<0.0001	1.58264
	Emotional neglect	0.38013	0.09814	3.87	0.0001	1.55632
Perceived stress	0.26737	0.07203	3.71	0.0002	2.26777	

Abbreviation: VIF=variance inflation factor.