

SUPPLEMENTARY MATERIAL

SUPPLEMENTARY TABLE S1. Antimicrobial susceptibility profiles of 17 F toxinotyping *C. perfringens* strains

Drug class	Antimicrobial agents	MIC50 µg/mL	MIC90 µg/mL	Range µg/mL	R%	I%	S%
Macrolides	Erythromycin	2	2	1 to >2	0.0	100.0	0.0
Fluoroquinolones	Ciprofloxacin	0.25	0.5	0.064 to 1	0.0	0.0	100.0
Lincosamides	Clindamycin	0.5	2	≤0.064 to 2	0.0	23.5	76.5
Oxazolidinones	Linezolid	1	1	0.5 to 1	0.0	0.0	100.0
Nitroimidazoles	Metronidazole	1	>32	0.5 to >32	47.1	0.0	52.9
Cephalosporins	Cefoxitin	0.5	4	0.064 to 1	0.0	0.0	100.0
Penicillins	Penicillin	≤0.064	1	≤0.064 to 1	0.0	23.5	76.5
	Amoxicillin	≤0.064	0.064	≤0.064	0.0	0.0	100.0
Tetracyclines	Tetracycline	1	4	≤0.064 to 16	5.8	5.9	88.2

Note: MIC90 and MIC50 values were defined as the lowest concentration of the antibiotic at which 90% and 50% of the isolates were inhibited, respectively.

Abbreviation: S=susceptible; I=intermediate resistant; R=resistant.